

New Probabilistic Strong Seismic Ground Motion map of Canada: a Compilation of Earthquake Source Zones, Methods and Results

P.W. Basham¹, D.H. Weichert², F.M. Anglin¹, and M.J. Berry¹.

Earth Physics Branch,
Energy, Mines and Resources Canada

1. Division of Seismology and Geomagnetism
Ottawa, Ontario K1A 0Y3

2. Pacific Geoscience Centre
Sidney, British Columbia V8L 4B2.

Earth Physics Branch Open File 82-33
Ottawa, Canada December, 1982.

205 pages including 67 figures

Price: \$61.50

Not for Reproduction

ABSTRACT

New probabilistic seismic ground motion maps of Canada, displaying peak horizontal acceleration and peak horizontal velocity at a probability of exceedence of 10 percent in 50 years, have been recommended as the replacement for the 1970 Seismic Zoning Map in National Building Code applications. This report presents a comprehensive description of the basic earthquake data and the methods employed in deriving the new maps.

RESUME

Les nouvelles cartes de probabilité des mouvements séismiques du sol pour le Canada ont été recommandées pour remplacer la carte de zonage séismique de 1970 dans les applications du Code national du bâtiment. Ces cartes présentent l'accélération horizontale maximum et la vitesse horizontale maximum à la probabilité de dépassement de 10 pourcent en 50 ans. Ce rapport donne une description détaillée des données séismiques de base ainsi que les méthodes utilisées dans l'élaboration de ces nouvelles cartes.

Table of Contents

	Page
ABSTRACT/RESUME-----	i
1. INTRODUCTION-----	1
2. CORNELL-MCGUIRE SEISMIC RISK ESTIMATION-----	6
2.1 Earthquake Source Zones-----	6
2.2 Magnitude Recurrence Relations-----	6
2.3 Strong Ground Motion Attenuation-----	12
2.4 Ground Motion Exceedence Computations-----	15
3. DESCRIPTION OF EARTHQUAKE SOURCE ZONES-----	15
3.1 Puget Sound (PGT)-----	15
3.2 Cascades (CAS)-----	19
3.3 Northern Vancouver Island (NVI)-----	23
3.4 Coast Mountains (CSM)-----	26
3.5 Juan de Fuca - Explorer (JFE)-----	29
3.6 Queen Charlotte Fault (QCF)-----	32
3.7 Sandspit (SPT)-----	35
3.8 Southeastern B.C. (SBC)-----	38
3.9 Flat Head Lake (FHL)-----	41
3.10 Northern B.C. (NBC)-----	41
3.11 Southern Saskatchewan (SAS)-----	45
3.12 Fairweather - Yakutat (FWY)-----	48
3.13 Denali - Shakwak (DSK)-----	52
3.14 Richardson Mountains (RIC)-----	55
3.15 Beaufort Sea (BFT)-----	58
3.16 Mackenzie (MKZ)-----	61
3.17 Alaska (ALC, ALI)-----	61
3.18 Charlevoix (CHV)-----	67
3.19 Western Quebec (WQU)-----	71
3.20 Lower St. Lawrence (LSL)-----	75
3.21 Northern Appalachians (NAP)-----	78
3.22 Laurentian Slope (LSP)-----	83
3.23 Attica (ATT)-----	86
3.24 Eastern Background (EBG)-----	90
3.25 Baffin Bay (BAB)-----	90
3.26 Baffin Island (BAI)-----	96
3.27 Labrador Sea (LAB)-----	99
3.28 Eastern Arctic Background (EAB)-----	102
3.29 Gustaf-Lougheed Arch (GLA)-----	105
3.30 Sverdrup (SVD)-----	109
3.31 Boothia-Ungava (BOU)-----	112

4.	PROBABILISTIC SEISMIC GROUND MOTION ANALYSIS-----	115
4.1	Regionalization-----	115
4.2	Integration over Source Zones-----	116
4.3	Treatment of other Errors-----	119
4.4	Limiting Ground Motion from Large Magnitudes-----	120
4.5	Statistical Scatter on Attenuation-----	120
4.6	Computational Grid and Contouring-----	120
4.7	Site-specific Risk Calculations-----	123
	ACKNOWLEDGEMENTS-----	123
	REFERENCES-----	125
	APPENDIX A-----	133
	APPENDIX B-----	161
	APPENDIX C-----	188

1. INTRODUCTION

In Canada, the primary application of seismic zoning information is made within the context of seismic loading provisions of the National Building Code (Associate Committee on the National Building Code, 1980). In the first edition of the code (1941) the seismic provisions appeared in an appendix and were based on concepts presented in the 1937 United States Uniform Building Code. In the 1953 edition, the earthquake loading requirements were updated and placed in the main text, and referenced the first seismic zoning map of Canada, which was subsequently described by Hodgson (1956). The Hodgson zoning map was a qualitative "seismic probability map" based on knowledge of the larger earthquakes and general considerations of the regional extent of earthquake zones.

The Hodgson zoning map was replaced in the 1970 edition of the code by the 1970 Seismic Zoning Map (Figure 1). This, the first strictly probabilistic map, was developed from the work of Milne and Davenport (1969) (see also Whitham et al., 1970), and displayed contours of peak horizontal acceleration at a probability of exceedence of 0.01 per annum that were used as boundaries for the four seismic risk zones. Although some of the seismic loading provisions have changed (Uzumeri et al., 1978), the 1970 zoning map has been referenced by subsequent editions of the code up to 1980.

The 1970 zoning map shown in Figure 1 was developed using extreme-value statistics applied to the catalog of known Canadian earthquakes (to 1963) to compute probabilities of peak acceleration exceedence at a grid of sites throughout the country (Milne and Davenport, 1969). Reviews and recent applications of seismic risk estimation in Canada (Weichert and Milne, 1979;




Figure 1

Basham and Weichert, 1979; Basham et al., 1979) have shown that the method developed by Cornell (1968) is the most appropriate for derivation of new probabilistic seismic ground motion maps of Canada. For computational purposes we have adapted the computer program of McGuire (1976) and will therefore refer to the method as "Cornell-McGuire".

The new probabilistic seismic ground motion maps are described in the publication by Basham et al. (1983), which includes a discussion and illustration of the influences of: (a) the expanded catalog of Canadian seismicity since the preparation of the 1970 map which was based on seismicity to 1963; (b) the change in method from extreme-value to Cornell-McGuire; (c) the change in strong ground motion attenuation relations from those of Milne and Davenport (1969) to the new relations developed by Hasegawa et al. (1981); and (d) a change in probability of exceedence from the value of 0.01 per annum used for the 1970 zoning map to the value of 10% in 50 years recommended for the next version of the National Building Code. The adaptation of probabilistic peak horizontal acceleration and velocity maps to seismic zoning maps and the concomitant changes to the seismic loading provisions that would be required in the National Building Code are described by Heidebrecht et al. (1983). The contour maps of acceleration and velocity at a probability of exceedence of 10% in 50 years that will be employed as new seismic zoning maps are shown in Figures 2 and 3.

It is the purpose of this report to present a comprehensive description of the basic earthquake data and the methods employed in deriving the new maps. The contents and format of the report are described in the following section with reference to the basic requirements of the Cornell-McGuire method.




Figure 2




Figure 3

2. CORNELL-MCGUIRE SEISMIC RISK ESTIMATION

The four basic components of the Cornell-McGuire seismic risk estimation method are illustrated schematically in Figure 4.

2.1 Earthquake Source Zones

The method requires that the seismicity be defined in finite source zones (Figure 4a) with uniform activity. The seismicity of Canada and adjacent active regions has been modelled with a total of 32 source zones (Figure 5) based on the distribution of historic and recent earthquakes and any geologic or tectonic evidence that can be employed to delineate the extent of future earthquake activity. A description of the rationale used for the selection of zone boundaries and a small scale map of each of the source zones, with its associated seismicity, is given in Section 3. The source zone boundaries on Figure 5 and on the individual source zone maps are straight lines in the Lambert Conformal projection used for these maps. Each of these zones is modelled as a horizontal, uniformly active source of earthquakes. In the absence of a reliable depth distribution, the seismicity in all zones, with one exception, is assigned a focal depth of 20 km. This is slightly deeper than the average depths of Canadian earthquakes, but the choice partially compensates for the unrestricted near-field attenuation (see Section 2.3). The exception is the Puget Sound subduction zone (see Section 3). The Alaskan seismicity is also modelled in simplified zones at a depth of 20 km, even though many of the earthquakes do occur in deeper subduction zones.

2.2 Magnitude Recurrence Relations

Each of the zones is assigned a cumulative magnitude recurrence relation




Figure 4




Figure 5

terminated at an upper-bound magnitude (Figure 4b). The recurrence relations have been computed using the method of Weichert (1980), a maximum likelihood method extended to the case of earthquakes with an assigned maximum magnitude and grouped in magnitude with each group observed over its individual time period. This requires an estimate of the first year of complete reporting of different magnitude category earthquakes in each of the zones. The estimated years for half-magnitude categories are given in Table 1. These are estimates based on our experience and on discussions with Branch colleagues familiar with historical Canadian seismicity, and are determined by the historical patterns of population distribution and reporting of earthquake occurrences in the pre-instrumental era, and by the capabilities of global and Canadian seismograph networks and methods of routinely reporting earthquakes that have developed since the turn of the century. Milne et al. (1978) and Rogers (1983) describe these considerations for a part of the west coast region (see also Basham and Whitham (1966)). For a number of zones a starting year is imposed rather arbitrarily on a larger magnitude category. These cases are noted in the individual zone descriptions in Section 3. The earthquakes that postdate these completeness years were used to derive the magnitude recurrence relations and are listed in Appendix A. The final year of earthquakes included for these computations is 1977, with two exceptions noted in Section 3 for which 1978 data were used. For only the Northern Appalachian zone would the inclusion of more recent earthquakes be expected to influence the derived magnitude recurrence relation; the implications for this zone are discussed in Section 3.21.

In the compilation of earthquakes for each of the source zones, aftershocks are included if they pass the completeness test described in Table 1. It is a debatable question whether aftershocks should be included in

Table 1

Estimated First Year of Complete Reporting of
Magnitude Categories

Zone	Magnitude Category*										
	3.0	3.5	4.0	4.5	5.0	5.5	6.0	6.5	7.0	7.5	8.0
<u>Western Canada</u>											
PGT	-	-	1965	1956	1940	1917	1899	1899	1860	1860	
CAS	-	-	1965	1956	1940	1917	1899	1899	1860	1860	
NVI	-	1965	1956	1956	1940	1917	1917	1917	1860	1860	
CSM	-	1965	1956	1956	1940	1917	1917	1917			
JFE	-	-	1965	1965	1965	1917	1917	1917	1899		
QCF	-	-	1965	1965	1965	1940	1917	1917	1899	1899	1899
SPT	-	-	1965	1965	1965	1940	1917	1917	1899		
SBC	-	1965	1960	1960	1940	1917	1899	1899			
NBC	-	1971	1965	1965	1965						
SAS	1968	1965	1940	1940	1940	1900					
<u>Northwestern</u>											
FWY	-	-	1972	1968	1964	1950	1950	1930	1920	1850	1850
DSK	-	-	1972	1968	1964	1950	1950	1930	1920		
RIC	-	-	1968	1968	1964	1950	1950	1930	1920		
BFT	-	-	1968	1968	1964	1950	1950	1930			
MKZ	-	-	1968	1968	1964	1950	1950				
<u>Eastern Canada</u>											
CHV	1968	1963	1937	1928	1920	1900	1800	1660	1660		
WQU	1968	1963	1937	1928	1928	1900	1900	1850			
LSL	1975	1963	1963	1937	1937	1937	1900				
NAP	1975	1963	1937	1937	1937	1900					
LSP	-	-	1956	1956	1937	1937	1930	1930	1800		
ATT	-	1963	1937	1937	1937	1850					
EBG	-	1963	1963	1956	1937						
<u>Northeastern</u>											
BAB	-	-	1968	1968	1964	1950	1950	1930	1920	1850	
BAI	-	-	1968	1968	1964	1950	1950				
LAB	-	-	1968	1968	1964	1950					
EAB	-	-	1968	1968	1964	1950					
GLA	-	-	1968	1968	1964	1950					
SVD	-	-	1968	1968	1964	1950					
BOU	-	-	1968	1968	1964	1950	1950				

*With magnitudes defined to one-tenth unit, each category includes earthquakes in a half-magnitude range; e.g., M 5.5 includes M 5.3 to 5.7.

defining magnitude recurrence relations for the purpose, here, of deriving earthquake source models for seismic risk estimates. On the one hand, the inclusion of aftershocks violates the assumption of Poissonian distribution often used to model earthquake occurrence; on the other, large aftershocks can contribute risk in their own right. Further, it is often difficult to decide if earthquakes have occurred as mainshock-aftershock sequences, or as swarms with many events of similar magnitude. Examples of swarm-like activity described in Section 3 are the earthquakes of Byam Martin Channel, Baffin Island and Miramichi, New Brunswick. In general, the effect on magnitude recurrence of including aftershocks is a small change in the recurrence slope. This may be a small increase if many small aftershocks pass the completeness test, or a small decrease if only large aftershocks of the larger historical earthquakes pass the completeness test.

Each of the magnitude recurrence relations is terminated by an adopted upper-bound magnitude. The upper bound magnitude truncates the incremental magnitude distribution which produces a smooth curve approach to zero rate in the plotted cumulative distribution.

The maximum magnitude earthquake that can occur in a source zone can be a critical parameter in probabilistic estimates of seismic risk. For zones with high rates of seismicity, significant risk contributions at moderate probabilities are coming from earthquakes near the maximum; therefore, the choice is important. However, for zones of low seismicity the probability of occurrence of earthquakes near the maximum can be much less than the probability being considered in the risk estimate, and the choice is less important.

There are a number of ways of estimating maximum magnitude: by a

magnitude truncation in observed seismicity for source zones in which the return period for maximum magnitude is shorter than the observation period; by consideration of the maximum fault area that can break in a single event; by estimates of the average fault slip rate, from plate tectonic models or geological data. Where this type of evidence is available, these methods are considered for choosing upper bound magnitudes. For many zones, however, this type of evidence is not available, and a rather arbitrary value has been adopted. In many cases this is approximately one-half a magnitude unit larger than the largest known historical event. A discussion is given in Section 3 if the choice is considered important to the resulting estimate of earthquake risk.

The magnitude recurrence relations have the form

$$N(>M) = N_0 \exp(-\beta M) (1 - \exp(-\beta (M_x - M))).$$

A summary of the recurrence parameters and the total area of each source zone is given in Table 2. Figures showing graphical illustrations of the recurrence curves accompany the zone descriptions in Section 3. In a number of cases the seismicity data are too sparse to derive an independent relation and recurrence parameters are imposed (parameters in parentheses in Table 2). For those source zones exclusively in United States territory we have not estimated the magnitude recurrence parameters but have adopted them from equivalent work by the U.S. Geological Survey.

2.3 Strong Ground Motion Attenuation

Attenuation relations that predict ground motion as a function of magnitude and distance (Figure 4c) are required for the ground motion parameters being mapped, and Hasegawa et al. (1981) have developed the

Table 2

Source Zone Magnitude Recurrence Parameters

Zone	β	N_0	M_x	Area (km ²)
<u>Western Canada</u>				
PGT	1.58	436.	7.5	28400
CAS	1.87	1060.	7.5	145000
NVI	1.04	21.	7.5	27000
CSM	1.77	272.	6.5	139000
JFE	1.72	7360.	7.0	84800
QCF	1.50	1610.	8.5	46400
SPT	1.87	1240.	7.0	53700
SBC	2.28	3230.	6.5	255000
NBC	(2.28)	(1830.)	5.0	875000
FHL	2.58	38000.	6.5	2100
SAS	2.07	188.	6.0	
<u>Northwestern</u>				
FWY	1.66	4590.	8.5	111000
DSK	1.96	2820.	7.0	110000
RIC	1.76	1560.	7.0	20000
BFT	1.76	681.	6.5	39000
MKZ	2.67	92000.	6.0	698000
ALC	1.43	3820.	8.5	132000
ALI	1.73	57100.	8.5	321000
<u>Eastern Canada</u>				
CHV	1.66	310.	7.5	6880
WQU	1.85	1030.	7.0	121000
LSL	1.85	533.	6.0	24500
NAP	1.87	638.	6.0	241000
LSP	1.30	41.	7.5	15200
ATT	1.32	11.	6.0	2620
EBG	2.78	16200.	5.0	2670000
<u>Northeastern</u>				
BAB	1.64	611.	7.5	100000
BAI	2.54	52100.	7.0	85000
LAB	1.95	1970.	6.5	352000
EAB	1.81	847.	6.0	1067000
GLA	2.19	18900.	6.5	42000
SVD	(2.19)	(2280.)	6.0	480000
BOU	2.02	3780.	6.5	830000

relations for this purpose in Canada. These authors, and Heidebrecht et al. (1983) and Basham et al. (1983), have discussed the need to estimate probabilistic ground motion in the two dominant frequency ranges represented by the parameters of peak horizontal acceleration (near 5 Hz) and peak horizontal velocity (near 1 Hz).

The analytical form of the attenuation relations of Hasegawa et al. (1981) are unrestricted at high magnitudes and in the near distances ranges. Although there are no strong motion data available for large earthquakes ($M \geq 7.5$) to provide good evidence, it is generally agreed that the excitation of seismic ground motion in the frequency range of engineering interest reaches an upper limit as magnitude increases to large values, much as the magnitude scales that measure ground motion in the frequency range near 1 Hz tend to saturate near M 7.5. To impose this condition on the Hasegawa et al. (1981) attenuation relations, the ground motion contributions from earthquakes with magnitudes greater than 7.5 are computed as if the earthquakes were magnitude 7.5; i.e., the recurrence relations for magnitudes greater than 7.5 are collapsed onto the relation for 7.5. The manner in which this is implemented in the computations is described in Section 4.4.

Although there is good evidence (e.g., Joyner and Boore, 1981; Campbell, 1981) that an extrapolation of attenuation relations applicable at greater distances to the near field will produce an over-estimate of peak ground motion parameters, particularly for the larger magnitudes, the Hasegawa et al. relations have not been explicitly restricted in the near field for purposes of these computations. There is, however, a de facto limitation on near field ground motion by the adoption of a minimum focal depth of 20 km for the earthquake source models. Hasegawa et al. (1982) have argued that this is

adequate for regional probabilistic ground motion mapping at moderate probabilities, but would not be adequate for estimating low-probability, near-field effects of large earthquakes that may be required for design of critical facilities (see also Basham et al. (1982) and Heidebrecht et al. (1983)).

2.4 Ground Motion Exceedence Computations

The final component in the Cornell-McGuire seismic risk analysis is the computation of a distribution function of probability of exceedence of the ground motion parameters (Figure 4d), by numerical integration of contributions from all relevant source zones. We have employed a modified version of the McGuire (1976) computer program; a program listing is given in Appendix B. A description of a variety of important matters related to the implementation of the program and the stability of the calculations is given in Section 4.

3. DESCRIPTION OF EARTHQUAKE SOURCE ZONES

3.1 Puget Sound (PGT) (Figures 6, 7)

The large Benioff zone of relatively deep earthquakes in this area is the most important factor differentiating this zone from the surrounding zones. The events appear to be in the subducted oceanic lithosphere, and may arise from the bend in the sinking oceanic lithosphere from a dip of 10-20° under the coast, to a dip of about 50° east of Puget Sound. Crosson (1982) has shown that diffuse shallow seismicity extends throughout this zone to depth of 20 - 25 km. A quiet zone separates this low-magnitude seismicity from deeper activity at 40 - 70 km, which dips towards the northwest. The





Figure 6



observed seismicity, when interpreted as seismic shortening, reflects only 10% of the convergence rate between the Juan de Fuca and North American plates in this area (Weichert and Hyndman, 1982a), but corresponds approximately to a north-south component of convergence, i.e., parallel to the margin. The corner of the continental margin (buried trench or axis of start of subduction) requires N-S compression or overlap in the subducted lithosphere in this area. Towards the north, subduction has slowed and is northward oblique to the margin (Hyndman et al. 1979a), but a detailed picture of the transition beneath the margin has not yet emerged.

A number of choices can be made in modelling the Puget Sound Zone. We have chosen to model the zone by a horizontal uniform distribution, at a depth of 40 km, of all events that occur within the zone boundaries. Although the more significant events may be deeper (40 - 60 km; Crosson (1982)), there is evidence (Hasegawa et al. 1981) that these events produce larger than average peak ground motion at epicentral distances smaller than their focal depths, which is partially accounted for by modelling them at the shallower depth. Crosson (1982) has shown that the Puget Sound seismicity rates are greater at smaller magnitudes in the shallow zone (0 - 30 km) and greater at larger magnitudes in the deep zone. Our model places all of the events at a depth of 40 km. This is an adequate simplification, but not strictly correct since the shallow seismicity described by Crosson has a higher activity than our Cascades zone (Section 3.2). Alternative models have been tested which include more of the Puget Sound activity in the overlapping and surrounding Cascades zone, but the risk estimates differ by only a few percent throughout the region.

The width of the Puget Sound zone is taken as about 50 km east and west of the estimated position of the change in dip of the subducted slab. The

eastern boundary of the zone is also set by calculations on the maximum possible landward persistence of Benioff-type events on thermal grounds. The landward persistence of oceanic lithosphere below the critical temperature for earthquakes depends on the rate of subduction and the age of the oceanic lithosphere being underthrust, both of which vary along the margin. The resulting model may be somewhat too wide in the east-west direction as the significant earthquakes tend to cluster near the centre of the zone (see Figure 6).

The northern boundary of the zone has been chosen along 49°N in agreement with the pattern of the larger historical events. Since a few smaller deep earthquakes have been observed a further 50 km north, under Georgia Strait, an alternate model would have to extend the zone that far. A third, more sophisticated model could include a gradual diminishing of activity and perhaps also of maximum magnitude from the south end of the Puget Sound to the north; our chosen model is thus intermediate in terms of the estimated risk to the densely populated Lower Mainland of British Columbia.

A maximum magnitude of 7.5 is selected for the zone as about half a magnitude unit larger than the largest in the data file, 7.1, in 1949. Magnitude 7.5 is approximately the earthquake size expected for a normal fault breaking completely through the subducted oceanic lithosphere (perhaps 20 km thick) over a horizontal dimension of 100 - 200 km, using the fault area - magnitude relation of Kanamori and Anderson (1975).

3.2 Cascades (CAS) (Figures 8, 9)

The shallower stress regime in this region probably arises from the Juan de Fuca - America plate convergence at a rate of several centimetres per





Figure 8



year. The eastern boundary of the zone is taken to the east side of the high Cascades on the assumption that their presence indicates a significant underlying change in stress regime. The eastern limit of the zone may also be taken from a probable eastward decrease in shear coupling between the continental lithosphere and the underthrusting oceanic lithosphere. The coupling and thus stress in the continental lithosphere may decrease as the temperature in the subducted oceanic lithosphere increases and the shear zone approaches the melting temperature under the volcanic zone.

Scattered seismicity extends from the coast to several hundred kilometres inland, and from southern Washington State to a quiet area in south-central Vancouver Island. The most significant earthquake in the historic record is the event of 1872 with an estimated magnitude somewhat greater than 7 (Coombs et al., 1976; Malone and Bor, 1979). The only obvious geologically-recent fault of a length that might generate such a large event seems to be the Fraser-Yalakom fault system although there is no evidence that the 1872 event occurred on this fault system. A maximum magnitude of 7.5 is selected to accommodate such an event anywhere in the zone, albeit at a rather low rate as shown by the magnitude recurrence curve (Figure 9).

In the Cascades zone model, and in all other zones described in the following, the earthquakes are assumed to occur at a focal depth of 20 km; i.e., the Puget Sound zone discussed above is the only one for which deeper focal depths are assumed. The Cascades zone is modelled to include the region above the Puget Sound Zone by assuming uniform shallow seismicity to extend throughout the area.

This overlapping of lower and higher seismicity zones occurs in a number of additional cases in the following. Although it is not the most

representative modelling of the seismicity that is possible, it is done to avoid the excessive calculations that would be required for these overlapping zones if they were modelled with a "cut-out" of the more active zones, which would require a more detailed pattern of sub-zones for the risk analysis.

3.3 Northern Vancouver Island (NVI) (Figures 10, 11)

The stress field in this area is related to the Explorer -Juan de Fuca-America plate interaction (Hyndman et al., 1979): varying rates of convergence along the margin and strike-slip across the offshore Nootka fault perpendicular to the margin. There may be stress coupling between the Nootka fault which is being subducted beneath the margin and the overlying continental lithosphere. The northern and southern limits of the zone are parallel to and roughly equidistant from the landward projection of the Nootka fault zone. Also included near the north end of the zone are the geologically recent plutons across Vancouver Island that probably arise from the northern edge of the subducted Explorer plate. The narrowing of the zone to the north is suggested by decreasing ocean-continent interaction as the oceanic plate becomes younger, thinner and weaker and the convergence rate becomes slower. The eastern boundary of the zone is taken as the edge of the Insular Belt.

As seen in air photos and satellite images, the zone exhibits some evidence of recent faulting in several areas. The most significant morphological feature, and the longest linear feature on the island, is the Beaufort Range scarp on which the magnitude 7.3, 1946 earthquake probably occurred (Rogers and Hasegawa, 1978; Slawson and Savage, 1979).

The seismicity includes a number of large shallow events, including the 1946 earthquake, in a roughly east-west line across north-central Vancouver





Figure 10



Island. There are, however, relatively few small events, which results in a low slope for the magnitude recurrence curve. In a somewhat arbitrary attempt to increase the slope, assuming the low value is in part due to a temporary lull in a numbers of small earthquakes in recent decades, we have included for this zone the 1978 earthquakes, most of which occurred near Cape Cook at the northwest corner of the zone. The inclusion of 1978 data increases the slope slightly but does not affect the estimated rates of the more significant higher magnitude events.

The maximum magnitude of 7.5 is selected as representative of a 100-km fault break with a depth of 20 km, i.e., a fault break with a length approximately half the largest dimension of the zone.

3.4 Coast Mountains (CSM) (Figures 12, 13)

A shallow stress regime can be postulated for this region primarily from the Explorer-America plate interaction along the margin, although the tectonic regime of the margin is complex. The eastern boundary is taken approximately at the eastern side of the Coast Mountains on the assumption that they reflect the limit of the major stress regime. The Coast Mountains zone can be considered as a lower level aureole around the Northern Vancouver zone, much like the Cascades zone around the Puget Sound zone, although the zone is modelled to overlap the Northern Vancouver Island zone.

The level of seismicity is quite low but is judged to be slightly higher than the adjacent Southeastern B.C. zone to the southeast (see Section 3.8). The northeast corner of the zone is chosen to include the historical events near Bella Coola that do not pass the completeness test; the western boundary terminates against the Juan de Fuca - Explorer zone. The maximum magnitude






Figure 12



of 6.5 is more than half a magnitude unit larger than the largest historic event, but is selected on the basis of assumed similarity with the Southeastern B.C. zone. However, neither the Cascades nor the Southeastern B.C. zones has known geologic or tectonic features that could be used to estimate maximum magnitude.

3.5 Juan de Fuca-Explorer (JFE) (Figures 14, 15)

The seismicity of this zone appears to follow the en-echelon ridge-transform boundary of the Pacific - Juan de Fuca plate boundary. (Riddihough, 1977; Hyndman et al. 1978; Riddihough et al., 1980; Davis and Riddihough, 1982). The Juan de Fuca ridge system consists of a series of spreading centres (Tuzo Wilson, Dellwood, Explorer, Juan de Fuca and Gorda), offset by transform fault segments (Dellwood-Wilson, Revere-Dellwood, Sovanco). The oceanic lithosphere landward of the Dellwood Wilson and Revere-Dellwood transform faults and Dellwood and Tuzo Wilson spreading centres appears to be coupled or nearly coupled to the America plate (Riddihough et al. 1980). Most of the seismicity is probably associated with transform faults rather than the ridge segments of the boundary. There is Plio-Pleistocene deformation and faulting in seismic profiles off the main plate boundaries nearer to the margin, such as the Winona ridge, but no clear evidence of more recent faulting. The zone is taken to extend from about 50 km west of the Pacific - Juan de Fuca plate boundary to the edge of the shelf, although the area of Winona Basin to the north of the Nootka fault zone appears to be less active. The width of the seismicity pattern probably comes from epicentral location uncertainties and biases. An ocean-bottom seismograph survey (Hyndman and Rogers, 1981) indicates that most of the active features are no more than 25 km wide.





Figure 14



The largest event in the data file is magnitude 6.7; there are many near this magnitude but none larger (Figure 15). There is good evidence for a geological limit on maximum magnitude. The faults on and near the plate boundary have a maximum length of about 100 km and could have a vertical extent of about 10 km; therefore the maximum fault area is about 1000 km². Using relations between magnitude and fault area (Kanamori and Andersen, 1975; Singh et al., 1980), the maximum plausible earthquake is magnitude 7.0. This is in good agreement with the historic data and is selected as the maximum magnitude (see also Hyndman and Weichert (1983)).

3.6 Queen Charlotte Fault (QCF) (Figures 16, 17)

The Queen Charlotte fault is the present transform boundary between the Pacific and North America lithospheric plates off western Canada between 52°N and 55°N. Off Queen Charlotte sound, there is a triple point with a convergence zone to the southeast and the Juan de Fuca ridge system to the southwest (Keen and Hyndman, 1979; Davis and Riddihough, 1982). The Queen Charlotte fault plate boundary has primarily right lateral, strike-slip motion with an average rate of about 55 mm per year (Atwater, 1970; Riddihough, 1977). Some convergence and underthrusting is predicted from global plate models (e.g., Minster and Jordan, 1978) and is also suggested by the shallow depression or trench and associated gravity low along the margin and by uplift over the Queen Charlotte Islands (e.g. Currie et al., 1980; Bird, 1981; Hyndman and Weichert, 1982b; Hyndman et al., 1982). The morphology and seismic profiles indicate two parallel fault scarps on the continental slope separated by a 30 km wide irregular terrace at a water depth of 2 km. The present seismic activity is concentrated on a probably vertical fault beneath the landward of the two slopes (Hyndman and Ellis, 1981).





Figure 16

ZONE: QUEEN CHARLOTTE FAULT (QCF)



The southern limit of the zone is taken as the triple point. The northern limit at 57°N, is a somewhat arbitrary division between the Queen Charlotte fault and the northern extension, the Fairweather fault system. The epicentres that fall west of the zone in Figure 16 may be mislocations of events that occurred along the fault; they are included in the zone for purposes of magnitude recurrence calculation.

The maximum magnitude selected for the Queen Charlotte Fault zone is 8.5. By integration of the magnitude recurrence relation (Figure 17), to provide an estimate of total seismic moment, Hyndman and Weichert (1983) estimate a slip rate of 52 mm per year on the Queen Charlotte fault system using a maximum magnitude of 8.5, i.e., in good agreement with the plate model estimate. From fault area considerations, this would represent a break along most of the length of the fault zone as defined in Figure 16.

3.7 Sandspit (SPT) (Figures 18, 19)

A number of recently active splinter faults trend generally northward to the east of the main Queen Charlotte fault zone, e.g., the Sandspit fault (Yorath and Chase, 1981; Yorath and Hyndman, 1983). Seismic profiling has revealed active grabens northeast of the Queen Charlotte Islands. The Rennell Sound-Louscoone Inlet fault is also a major fault trace. These faults could be a response to the small difference between the estimated direction of the Pacific-North America relative plate motion and the strike of the Queen Charlotte fault along the margin and the postulated very oblique underthrusting. The Sandspit zone covers these faults but does not extend as far east as the mainland coast. Its length and width are arbitrarily chosen to be the same as the Queen Charlotte Fault zone.





Figure 18



EARTH PHYSICS BRANCH EMR OTTAWA CANADA
DIRECTION DE LA PHYSIQUE DU GLOBE OTTAWA CANADA

01/10/81 15.41.58.

Figure 19

The seismicity of the Sandspit zone proper is difficult to distinguish from apparent seismicity that may be caused by mis-location of Queen Charlotte fault earthquakes. We included only those events that are not obvious mis-locations; however, there is no certainty that all of the events actually occurred in this zone, or that there are not some mis-located Sandspit zone events included with the Queen Charlotte Fault zone seismicity.

Rogers (1982) has subsequently reviewed the older events and concluded that most of the seismicity that we have included in this zone occurred on the Queen Charlotte fault. Our source zone models were finalized prior to Rogers' work and, therefore, his revisions have not been included. The effect on the probabilistic ground motion maps are, however, negligible because the overwhelmingly dominant source zone in this region is the Queen Charlotte fault. There have also been recent (1982) earthquakes located in the Sandspit zone and east of it in Hecate Strait, so the Sandspit zone has been retained.

3.8 Southeastern B.C. (SBC) (Figures 20, 21)

All of the interior of B.C. could be considered a typical background zone. However, the differences in detection completeness between north and south and an apparent higher seismicity in the south are reasons for considering a separate southeastern zone. In the Rocky Mountain part of the zone the high topography may reflect higher than average stress, but tectonically the Rocky Mountain area is probably not related to the Intermountain seismic belt to the south. A hot spot trace similar to, but weaker than the Yellowstone hot spot, may traverse British Columbia west to east with the youngest rock ages lying near the 1918 magnitude 6 earthquake north of Revelstoke (Rogers and Ellis, 1979; Rogers et al., 1980; Rogers, 1981). We consider this centre of activity not yet sufficiently well defined





Figure 20

ZONE: SOUTHEASTERN B.C. (SBC)

CUMULATIVE RATE (P.A.)




to justify a separate zone. The changing epicentral pattern in the 1960s and 1970s is evidence against the notion of a quiet zone between the Rockies and the Cascades zone. The west-central portion of the zone as displayed in Figure 20 is devoid of earthquakes, but this region has experienced earthquakes as large as M 4 that do not pass the completeness criterion for inclusion here.

3.9 Flat Head Lake (FHL) (Figures 20, 22)

This zone is part of the Intermontane Seismic Belt, extending north-south in west-central United States as far north as Flathead Lake, Montana. The belt has been interpreted as a boundary within the main North America lithospheric plate. Two subplates are moving apart producing rift faulting (e.g. Smith and Sbar, 1974). The seismic activity is characterized by shallow focal depths and swarm activity. The nearest concentration to Canada is near 48°N at Flathead Lake, and we consider this to be the only significant contribution to Canadian seismic risk. We have modelled the source with a small area (shown on Figure 20 with the Southeastern B.C. zone) with activity scaled to match source 27 of Algermissen and Perkins (1976). Instead of their maximum magnitude of 5.5, we have adopted 6.5, which is still about 1/2 unit smaller than the maximum observed in the Intermontane seismic belt (i.e. Hebgen Lake, 1959). The resulting magnitude recurrence curve is shown in Figure 22.

3.10 Northern B.C. (NBC) (Figures 23, 24)

This is a zone of very low seismicity that includes the northern B.C. Cordillera. It has been extended into the Yukon to include the region between







Figure 23



EARTH PHYSICS BRANCH EMR OTTAWA CANADA
DIRECTION DE LA PHYSIQUE DU GLOBE OTTAWA CANADA

30/10/81 10.39.43.

Figure 24

the Shakkwak Valley and the Tintina Trench (see Section 3.3). The detection threshold at low magnitudes in this zone has been poor until very recently. (Three new regional stations were installed in 1981.) However there are no known events as large as magnitude 5, which would have been detected since at least the early 1960's. Tectonically, this zone cannot be considered to be much different from southeastern B.C., and the magnitude recurrence slope for the later zone has been imposed on the meagre data base to produce the recurrence relation shown in Figure 24. The maximum magnitude has been set at 5.0, one-half unit above the observed maximum. However, with the low earthquake rates in a relatively large zone its contribution to the probabilistic ground motion will be small.

3.11 Southern Saskatchewan (SAS) Figures 25, 26)

This source zone has been drawn to encompass the cluster of seismicity in southern Saskatchewan and adjacent Montana and North Dakota (Horner and Hasegawa, 1978). The main cluster of earthquakes, including the M 5.5 event in 1909, is spatially associated with the Williston Basin, but the zone is extended to the southwest in Montana to include magnitude 3-4 earthquakes that occurred in the 1969-1973 time period. The location of the 1909 earthquake is not well known, but the location of 49°N, 104°W was selected by Horner and Hasegawa (1978) as the centre of the area of maximum intensity.

There is good evidence (Horner et al., 1973; McLennan et al., 1983) that the earthquakes included in the Southern Saskatchewan source zone are tectonic events in the Precambrian basement. There is also evidence (Gendzwil et al., 1982) of earthquakes as large as M 3.5 as far north as central Saskatchewan being induced by potash mining activity. These induced events have produced






Figure 25

ZONE: SOUTHERN SASKATCHEWAN (SAS)



EARTH PHYSICS BRANCH EMR OTTAWA CANADA
DIRECTION DE LA PHYSIQUE DU GLOBE OTTAWA CANADA

29/10/81 16.09.48.

Figure 26

moderate seismic shaking, but they cannot be included in a general model of earthquake sources because regions of similar potash mining in future are not known.

The maximum magnitude chosen is 6.0, half a unit larger than the 1909 earthquake, but the adopted recurrence relation is poorly defined (Figure 26). This source zone contributes a small region of peak acceleration greater than the minimum level contoured (Figure 2), but does not contribute significantly to peak velocity (Figure 3).

3.12 Fairweather-Yakutat (FWY) (Figures 27, 28).

The region of transitional tectonics, from transcurrent faulting along the coast of British Columbia and southeastern Alaska to subduction along the Aleutian Island arc (e.g., Perez and Jacob, 1980; von Heune et al., 1979) is modelled here as one continuous, simplified zone (Figure 27). The southern boundary at 57°N is the approximate location of the transition from the Queen Charlotte Fault offshore to the Fairweather Fault onshore in southeastern Alaska. The northeastern and northern boundaries are drawn to include in the zone the large earthquakes of the strike-slip Fairweather and underthrusting Chugach - St. Elias Faults, respectively. The western boundary is somewhat arbitrary, but it is drawn at 145°W which is the easternmost extent of the rupture zone of the great Alaska earthquake of 1964 and, in the offshore, is the approximate location of the beginning of the Aleutian Trench. (The seismicity to the west is modelled as separate zones described below.) The southern boundary is drawn along the shelf edge structure to include in the zone the seismicity of the Yakutat Block.

This source zone includes the series of large (magnitude 8) earthquakes





Figure 27

ZONE: FAIRWEATHER-YAKUTAT (FWY)



which ruptured the coast between Yakutat Bay and Kayak Island at the turn of the century, and the magnitude 7.9 on the Fairweather fault in 1958. The inclusion of the turn-of-the-century earthquakes for the magnitude recurrence estimates seems to produce too high a rate for the larger earthquakes compared to a well defined relation for events less than magnitude 7 (Figure 28). To partially reduce this effect the completeness date for earthquakes of magnitude 7.5 and greater has been extended back to 1850 (Table 1), even though we don't believe there has been complete reporting of even these larger events since that date. The reduction of the starting year to 1850 simply imposes an assumption that no large earthquakes occurred in the zone between 1850 and 1899.

On the other hand, this region has been identified as a seismic gap. Sykes (1971) identified a gap between the aftershock zone of the 1964 Prince William Sound and the 1958 Fairweather Fault earthquakes. Lahr et al. (1980) demonstrated that this gap was only partially filled by the 1979 St. Elias earthquake. If the Pacific and North American plates have been converging at the rate of 5 cm/yr since the turn of the century, enough elastic strain has accumulated to produce a potential slip of 4 m. If this amount of slip occurred in one earthquake, it would generate an event as large as magnitude 8 that would likely fill the remainder of the gap (Lahr and Plafker, 1980).

The influence of potential seismic gaps is not included in this probabilistic analysis. It is sufficient to note here that the seismic ground motion on Canadian territory in the southwestern Yukon (we are not attempting to predict ground motion on U.S. territory in which most of this zone is located) at moderate probabilities is dominated by the large earthquakes in the zone (see Figure 28). In essence, the analysis includes the effects of a

gap-filling earthquake, because the model assumes that an earthquake near the maximum magnitude can occur with equal probability anywhere in the zone.

A maximum magnitude of 8.5 has been used for this zone, on grounds similar to those described above for the Queen Charlotte Fault zone. However, in this case the largest earthquake could be either primarily strike-slip on the Fairweather fault system, or primarily underthrusting on the Chugach-St. Elias fault system.

3.13 Denali-Shakwak (DSK) (Figures 29, 30)

This zone includes the seismicity between the Fairweather-Yakutat zone and the Denali-Shakwak-Dalton fault system. Its northeast edge includes this fault system, but the area further northeast to the Tintina Trench is relatively aseismic and has been included with the Northern B.C. zone described above. The easternmost boundary is drawn to include the inferred faulting along Chatham Strait (see, e.g., Figure 1 in Perez and Jacob (1980)), although recent results (Horner, 1983) have shown Chatham Strait to be essentially aseismic at low levels, with the seismicity trending southward toward the coast through the region of Glacier Bay. As with the Fairweather-Yakutat zone, the western boundary is selected as 145°W, with the seismicity further west in Alaska included in separate zones described below.

Horner (1983) has shown that both the larger historical events and the low level seismicity in recent years is quite restricted, west of the Alaska-Yukon border, to a narrow zone following the Duke River, Shakwak and Dalton Fault zones, i.e., the seismicity is likely confined to a narrower zone along known faults than the source zone employed here (Figure 29). The largest known historic event was magnitude 6.5 in 1944 near Haines Junction.





Figure 29



The upper-bound magnitude of 7.0 selected for this zone carries the assumption that larger events typical of major plate interactions in the Fairweather-Yakutat zone will not occur. However, the tectonics of the zone, which must bear some relation to the plate interactions along the margin of the Gulf of Alaska, is not well-understood.

3.14 Richardson Mountains (RIC) (Figures 31, 32)

This zone is a relatively confined, but highly active region of the northeastern Yukon. Recent reassessment of the locations of the larger pre-1966 events has enabled the dominant seismicity to be enclosed by a zone about 80 by 250 km (Figure 31). There is evidence, however, (Leblanc and Wetmiller, 1974) that the zone may be made up of two clusters, one which is centred in the Richardson Mountains and another in the Mackenzie Mountains, with a relatively quiet region between the two in the area of the Bonnet Plume Basin.

The tectonic process responsible for this cluster of seismicity, and any relationships between it and major plate interactions in the Gulf of Alaska or the opening of the Arctic Ocean basin are not yet understood. The only evidence to associate seismicity with local geological features (Norris, 1972) is given by Leblanc and Wetmiller (1974). They show a spatial correlation of low level activity detected in a 1972 field experiment with mapped extensional and transcurrent faults. A single, but poorly controlled, P-nodal solution for a magnitude 4.8 earthquake shows right-lateral motion on a nearly vertical fault with the same strike as the mapped faults.

The largest known historic event was magnitude 6.6 in 1955; the upper-bound magnitude is selected as 7.0.





Figure 31



3.15 Beaufort Sea (BFT) (Figures 33, 34)

The cluster of seismicity in the Beaufort Sea is another example of relatively confined cluster with poorly-understood tectonic cause. The zone boundaries in Figure 33 have been drawn to enclose the distribution of activity that has been shown by joint-epicentre solutions (Hasegawa et al., 1979) to be real, and not due to mislocations of previously catalogued earthquakes. The largest historic event, magnitude about 6.5 in 1920, does not pass the test for completeness, set at 1930, and so is not included in Figure 33 or Appendix A. However, the northeastern corner of the zone has been drawn to enclose the best available location for this event derived by Basham et al. (1977).

The earthquakes are confined to the region beneath the continental slope, between the 200 and 2400 m bathymetry contours, and fall between the seaward -20 mGal and landward +40 mGal contours of an elliptically shaped free-air gravity anomaly. Hasegawa et al. (1979) derived focal parameters for a 1975, magnitude 5.1 earthquake which suggested strike-slip motion on a steeply dipping fault plane at a depth of 40 km. The depth is unusual, as it may place the earthquake in the upper mantle beneath the continental margin. It is supported, however, by the hypothesis of high horizontal deviatoric stresses due to an uncompensated load of Quaternary sediments, which would produce maximum stress at approximately this depth. This, or some other stress, is acting on unhealed faults at, at least, lower crustal depths to produce the cluster of Beaufort Sea seismicity, but there is no geological or geophysical evidence to determine the real nature of the faulting beneath the continental slope.

There is only the one known earthquake in 1920 with magnitude of about 6





Figure 33



and two with magnitude 5, in 1937 and 1975. The upper-bound magnitude has been set at 6.5, but the rates of the larger earthquakes and the maximum magnitude must be considered rather poorly defined (Figure 34).

3.16 Mackenzie (MKZ) (Figures 35, 36)

This is a zone of "background" seismicity in the western Yukon Cordillera, surrounding the active Richardson Mountains zone and abutting the Beaufort Sea zone. It is bounded on the southwest by the Tintina Trench, on the southeast by the physiographic limit of the Cordillera in the region of the Liard River, and on the northeast by the Mackenzie River.

The seismicity includes the swarm of earthquakes off Martin Point, Alaska, with magnitudes as large as 5.3, most of which occurred in 1968, and scattered events east of the Mackenzie Delta and throughout the Yukon-Northwest Territories border region. Basham et al. (1977; their Figure 11) suggest that this seismicity, like that in the Richardson Mountains zone, is spatially correlated with the areas of most severe geologically mapped faulting.

The upper-bound magnitude has been set a 6.0 but, even though there is a relatively large number of earthquakes in the zone, the rates of the larger events is poorly defined (Figure 36).

3.17 Alaska (ALC, ALI) (Figures 37, 38, 39)

Thenhaus et al. (1979) have defined 24 separate earthquake source zones for purposes of estimating probabilistic seismic ground motion in the region of Alaska and the adjacent continental shelves. For our purpose, of estimating the contributions of Alaskan earthquakes to seismic ground motion






Figure 35



EARTH PHYSICS BRANCH EMR OTTAWA CANADA


05/10/81 14.44.59.

DIRECTION DE LA PHYSIQUE DU GLOBE OTTAWA CANADA Figure 36




EARTH PHYSICS BRANCH
DIRECTION DE LA PHYSIQUE DU GLOBE EMR OTTAWA CANADA

Figure 37



ZONE: INLAND ALASKA (ALI)



EARTH PHYSICS BRANCH EMR OTTAWA CANADA
 DIRECTION DE LA PHYSIQUE DU GLOBE OTTAWA CANADA

30/10/81 10.39.43.


Figure 39

on Canadian territory, these source zones can be greatly simplified. For the region of Alaska west of 145°W the Thenhaus et al. source zones have been combined into two zones, Coastal Alaska (ALC) and Inland Alaska (ALI), (Figure 37). The ALC zone is essentially their zone number 23; the ALI zone a combination of seven of their zones, numbers 8, 14, 15, 16, 21, 22 and 24. The magnitude recurrence relation for ALC (Figure 38) is adopted directly from their zone 23; the relation for ALI (Figure 39) is the sum of the individual recurrence rates for their seven zones.

These zones have been extended only to 160°W as even large earthquakes further west in the Aleutian Islands will make negligible contributions to ground motion in Canada. Thenhaus et al. have derived minor source zones for northern Alaska, but this seismicity is adequately represented by the Mackenzie zone which extends to 145°W (Figure 35).


3.18 Charlevoix (CHV) (Figures 40, 41)

The Charlevoix zone is historically the most active zone in eastern Canada with at least five earthquakes with magnitude of 6 or greater (1663, 1791, 1860, 1870 and 1925). The 1925 event is the only earthquake with magnitude near 7 on land in eastern North America in the twentieth century. As part of the review of eastern Canadian seismicity by Basham et al. (1979), the magnitudes of a number of Charlevoix and other earthquakes were revised as listed in their Table 1. To provide the information necessary to make equivalent changes to the master Canadian Earthquake Epicentre File, a documentation of the revised parameters is included here as Appendix C. It should be noted that these revisions are based on a less than exhaustive



EARTH PHYSICS BRANCH
DIRECTION DE LA PHYSIQUE DU GLOBE EMR OTTAWA CANADA

Figure 40



reassessment of all available data, but they are considered improvements on the parameters originally determined by Smith (1962, 1966).

The Charlevoix zone is interpreted as a region of steeply-dipping rift faults at the Shield-Paleozoic contact that has been weakened by a Late Devonian meteorite impact (Rondot, 1979; Hasegawa and Wetmiller, 1980; Anglin and Buchbinder, 1981; Basham et al., 1982). A projection of microearthquake activity to the surface along the postulated faults suggests that the active zone is confined between mapped faults on the north shore and a bathymetric feature in, and parallel to, the river near the south shore (Berry et al., 1982). Focal mechanism solutions (Leblanc and Buchbinder, 1977; Hasegawa and Wetmiller, 1980) indicate that high horizontal compressive stresses are now producing thrusting on the preexisting faults (Hasegawa and Adams, 1981).

The source model employed (Figure 40) is based on the distribution of historical seismicity. It is recognized that some, if not all, of the epicentres in the northwestern portion of the zone may be mislocations of events that occurred in the more confined source centred along the river (see Figure 12 of Basham et al. (1982)). Stevens (1980) has demonstrated that the larger events in the twentieth century, previously located elsewhere, had epicentres at either end of the confined zone; but data are not available to demonstrate this conclusively for the older events. Most of the larger, pre-instrumental earthquakes have been assigned locations in or near the river on the basis of macroseismic effects, but these may be biased because much of the early settlement was along the river.

In any case, the effect on the probabilistic ground motion results of the choice between the more confined or the larger historical source zone is

negligible away from the immediate vicinity of the zone. Near, or within, the zone the probabilistic results give little more than an indication of high earthquake risk; design considerations would be based on a more rigorous assessment of the expected near-field effects of large earthquakes.

The active zone described above has a length of about 80 km and the microearthquake activity suggests a depth of about 20 km. If a fault system the length and depth of the zone ruptured in one earthquake, it would have a magnitude of about 7.5 (Basham et al., 1982). This has been adopted as the maximum magnitude.

3.19 Western Quebec (WQU) (Figures 42, 43)

The boundaries of the Western Quebec zone (Figure 42) have been drawn to enclose a significant cluster of Shield earthquakes, the tectonic causes of which have been the subject of considerable research in recent years, but which remain poorly understood (Basham et al., 1979; Forsyth, 1981; Hasegawa and Adams, 1981; Forsyth et al., 1982).

The greatest number of earthquakes in this zone in recent years, although none with magnitude greater than 4.2, have been located in the central portion of the zone in Quebec north of the Ottawa river. Historically, the larger earthquakes have occurred on the fringes and outside of this concentration of recent events. An earthquake with magnitude about 6 occurred at or near Montreal in 1732. The magnitude of this event has been reduced to 6, from the previous 7, on the Canadian Earthquake Epicentre File on the basis of Leblanc's (1981) reassessment of the macroseismic data. During the twentieth century, earthquakes of magnitude 6.2 occurred near





Figure 42



EARTH PHYSICS BRANCH EMR OTTAWA CANADA
DIRECTION DE LA PHYSIQUE DU GLOBE OTTAWA CANADA

29/10/81 10.21.08.

Figure 43

Timiskaming in 1935 and magnitude 5.6 near Cornwall in 1944. The northern boundary of the zone is controlled by two earthquakes: a magnitude 5 that occurred the day following the Timiskaming event, but which Smith (1966) was convinced (by the available instrumental data) was at a different location; and a magnitude 5 event in 1950 near the headwaters of the Gatineau river. Both of these earthquakes are outside of the cluster of recent activity (see Figure 42). The southern boundary of the zone is extended into the Adirondacks and Lake Champlain region of New York State and Vermont, which has experienced similar low level activity but no large historic events.

Forsyth (1981) has shown that most of the earthquakes of western Quebec are located near or within the boundaries of the northeastern part of the Grenville metasedimentary belt and near the junction of the rift structures following the northern and eastern segments of the Ottawa river, the St. Lawrence river and Lake Champlain. The larger historic earthquakes (Montreal, Timiskaming, Cornwall) are spatially associated with these younger rift zones. The geological and aeromagnetic data indicate a Precambrian shear zone is continuous along most of the eastern side of the belt. The aeromagnetic and gravity data show distinct anomalies that suggest unmapped features along the northwest side. The seismicity in the central portion of the zone coincides with the interval between two prominent anomalies in the smoothed Bouguer gravity field, and shows a spatial correlation with a topographic regional low.

It appears that the seismicity reflects adjustment to a stress field resulting from one or more of: regional density variations, continental deglaciation and intraplate forces (see also Hasegawa and Adams (1981)).

However, the relative effect of each stress field and the reason for greater recent seismicity in the Grenville metasedimentary belt remains unclear.

Forsyth et al. (1982) have recently extended these and Landsat lineament correlations northward toward the Kapuskasing Fault Zone. They also show that the Kapuskasing region has been more active than the Timiskaming region in recent years. However, for our purposes the Kapuskasing region has not been attached to the Western Quebec zone; it remains in the more diffuse Eastern Background zone discussed below.

A maximum magnitude of 7.0 has been chosen for the Western Quebec zone, but there is no seismological or geological evidence that we can employ to support this, or to demonstrate that some larger value may not be more appropriate.

3.20 Lower St. Lawrence (LSL) (Figure 44, 45).

The Lower St. Lawrence zone is a cluster of seismicity centred approximately over the north shore of the river in the region from Baie Comeau to Sept-Iles. The better-located events in recent years have epicentres in the river and on the north shore (Figure 44); some of the older events that have epicentres on the south shore (northern portion of Gaspé Peninsula) may be mislocations due to poor network control. The small number of earthquakes, some quite recent, that have occurred in Jacques Cartier Passage and on the adjacent north shore are not included in this zone, but in the Eastern Background zone (see Figure 52).

The magnitude 4.1 earthquake that was induced by the filling of the Manic 3 reservoir (Leblanc and Anglin, 1978) is not included in this data set. The northwestern corner of the zone boundary has been confined to the





Figure 44

ZONE: LOWER ST. LAWRENCE (LSL)

CUMULATIVE RATE (P.A.)



EARTH PHYSICS BRANCH EMR OTTAWA CANADA
DIRECTION DE LA PHYSIQUE DU GLOBE OTTAWA CANADA

29/10/81 10.21.08.

Figure 45

south of the Manic 3 dam. However, the small earthquakes that are included in the northwestern corner of the zone, most of which occurred in 1966, may have been related to the 1965 filling of the Manic 2 reservoir, but failed to be recognized as such (see Figures 1 and 2 of Leblanc and Anglin). Because of the uncertainty, they are assumed here to be natural tectonic events.

Geological and geophysical features that may correlate with and control the Lower St. Lawrence zone are much less well known than is the case for the Charlevoix and Western Quebec zones. Among the sparse evidence is the study by Goodacre and Hasegawa (1980) showing that earthquakes in the Quebec City to Sept-Iles region of the St. Lawrence valley tend to cluster in regions of negative free-air gravity anomalies that are adjacent to major free-air gravity highs. In the region of the Lower St. Lawrence zone there is a small gravity high south of Sept-Iles, with the St. Lawrence river negative anomaly on each side, and a positive anomaly on the northern portion of the Gaspé peninsula that is part of the linear belt of positive anomalies south of the Appalachian front. Goodacre and Hasegawa suggest that gravitationally induced stresses, superimposed on an ambient tectonic stress field may be sufficient to activate pre-existing faults.

3.21 Northern Appalachians (NAP) (Figures 46, 47)

The Northern Appalachians zone is a relatively large zone of rather uniform seismicity throughout New Brunswick, Maine, New Hampshire and Vermont. The southeastern boundary of the zone is drawn to include the seismicity in the Bay of Fundy and off the coast of Maine. The southern boundary is an arbitrary one, adopted by Basham et al. (1979), that excludes from consideration the seismicity in southern New Hampshire and Massachusetts,




Figure 46




Figure 47

in particular the large number of historic events that are catalogued for the Boston region during the time of early settlement. The zone is extended to the southeast far enough to include the seismicity of southeastern New York along the Hudson River, but it is terminated north of the Ramapo fault system shown by Aggarwal and Sykes (1978) and Yang and Aggarwal (1981) to be active at low magnitudes. The northern boundary of the zone is drawn along the southern edge of the 100- to 200-km wide, relatively aseismic band that extends from the eastern side of Lake Champlain to the Gaspé Peninsula. Thus, the boundaries of this zone in the United States are arbitrary ones that are not intended to bear a particular relation to the structural geology and tectonics; the Canadian boundaries distinguish the zone from the relatively aseismic areas that include the Gulf of St. Lawrence, P.E.I. and Nova Scotia.

The structural grain of the Northern Appalachians is controlled by northeasterly trending belts of volcanic and sedimentary rocks of Devonian to Ordovician age that are intruded by post-Ordovician granites and basic dykes. Wetmiller (1975) found one plane of the focal mechanism of the 1973, M4.8 Quebec-Maine border earthquake to be on strike with the regional Appalachian trend. However, Yang and Aggarwal (1981) determined thrust faulting on a north-striking plane for this earthquake, and high-angle reverse faulting on north- to northeast-striking planes for 12 other earthquakes along the eastern margin of the Appalachians. The results available at the time of writing for the 1982 Miramichi, New Brunswick earthquakes (Wetmiller et al., 1982; Stevens, 1982) also suggest thrust faulting on north-striking planes. The Miramichi earthquakes, however, present an excellent example of the difficulties of making a clear correlation between even exceptionally well-documented and shallow earthquakes and the local geological features, a difficulty that, no doubt, pertains to much of the Northern Appalachian zone.

The larger historic earthquakes in the Northern Appalachian zone, as defined in Figure 46, have magnitudes estimated as about 5. These include the 1869 and 1904 events that caused minor damage in southern New Brunswick and eastern Maine, the 1940 events near Ossipee Lake, New Hampshire, the 1943 event near Dover-Foxcraft, Maine, and the 1973 Quebec-Maine border event. On the basis of these events, the maximum magnitude adopted for the recurrence relation is 6.0 (Figure 47). There is no geological or seismological evidence on which to base the maximum magnitude.

The Miramichi earthquake sequence of 1982 is an unprecedented sequence for eastern Canada, although the larger events, M5.7 and 5.4, are considered typical of the more significant earthquakes that can occur in the Northern Appalachian zone. To illustrate the influence this sequence has on the magnitude recurrence relation adopted for the zone, the earthquakes have been updated to mid-1982 (listed in Appendix A) and the magnitude recurrence relation recomputed as shown by the open circles and dashed curve in Figure 47. The large numbers of small earthquakes in the Miramichi aftershock sequence increases the slope of the recurrence relation. However, with the maximum magnitude kept at 6.0, the addition of the Miramichi events does not significantly affect the estimated rates at larger magnitudes. In fact, the recent events provide an estimate of the rate at magnitude 5.5 that agrees very well with the extrapolation based on the pre-1978 events. The updated recurrence relation would increase the Northern Appalachian probabilistic ground motion only slightly (about 5 percent) and would not significantly change the contour patterns of Figures 2 and 3.

3.22 Laurentian Slope (LSP) (Figures 48, 49)

The Laurentian Slope zone is a small cluster of earthquakes at the edge of the continental slope at the mouth of the Laurentian Channel that includes one major event, the magnitude 7.2 earthquake of 1929 (Doxsee, 1948), one of only two magnitude 7 earthquakes known to have occurred offshore of eastern Canada.

Preliminary examination of available data suggests that the zone of earthquakes is spatially distributed approximately as seen in Figure 48; i.e., the scatter is not due to mislocations of events that occurred at or near one epicentre (Basham and Adams, 1982). The zone boundaries drawn to enclose the cluster are controlled on the east and west by the margins of the Laurentian Channel, on the north by the faults associated with the Orpheus Graben, and on the south approximately at the base of the continental slope (see Figure 3 of Basham and Adams). King (1979) has suggested that the earthquakes appear to be associated with the Glooscap fault, the combined Cobequid-Chetabucto-Orpheus Graben-Laurentian Channel fault system. The seismic reflection profiling used to locate the faults in the region of the channel indicate that most, and perhaps all, of the offset is pre-Pleistocene, but there is not sufficient resolution in the profiling to detect recent offsets if they were present in the youngest sediments. A study of aerial photography of this fault system where it crosses Nova Scotia has shown no evidence of fault linears or scarps in surface deposits that would suggest recent movement (D.R. Grant, personal communication, 1982).

The rate of 1929-sized earthquakes is poorly determined by the magnitude recurrence data. For purposes of computing Figure 49 it has been assumed that magnitude 7 earthquakes would have been completely reported since 1800 (see




Figure 48

ZONE: LAURENTIAN SLOPE (LSP)




Table 1), although we are by no means certain that reports of effects would even approximately locate such an offshore event in the early 1800's, had one occurred. Nevertheless, the result is a recurrence relation that shows reasonable agreement between the rates of larger earthquakes and the rates in the magnitude 4-5 range, although the slope of the curve tends to be lower than that of most other source zones.

This source zone produces a small region of high amplitude ground motion on the probabilistic maps (Figures 2 and 3), similar to that produced by the Charlevoix zone. Implicit in the adoption of this model is the assumption that the next large earthquake in the region will occur within the restricted zone at the mouth of the Laurentian Channel, i.e., rather than at some other location on the Newfoundland or Scotian Shelf. The evidence to support this assumption is not very strong, but we consider the model to be the best available for the present purposes. The result, however, is that the remainder of the Newfoundland and Scotia shelves falls within a zone of low background seismicity (Figure 52), which may under-estimate the real risk in these regions.

3.23 Attica (ATT) (Figures 50, 51)

The Attica zone has been drawn to enclose the M 5.5, 1929 Attica earthquake and the smaller M 3.5 - 4.5 events that have occurred near the south shore of Lake Ontario. Basham et al. (1979) defined this zone as extending through the Niagara Peninsula to the Hamilton-Burlington region. We now believe that many of the small earthquakes around the western end of Lake Ontario are the result of shallow pop-up phenomena, and have included this area in the general Eastern Background zone.





Figure 50



EARTH PHYSICS BRANCH EMR OTTAWA CANADA
DIRECTION DE LA PHYSIQUE DU GLOBE OTTAWA CANADA

29/10/81 10.21.08.

Figure 51

The most significant tectonic feature in the region is the north-south trending Clarendon-Linden structure that extends for over 100 km from Lake Ontario to the New York - Pennsylvania border (Fletcher and Sykes, 1977). There is no conclusive evidence that the 1929 earthquake occurred on this structure, but Fletcher and Sykes have shown seismicity induced by hydraulic mining activity to have thrust mechanisms on a plane nearly parallel to the Clarendon-Linden fault, and that other nearby natural events may be associated with branches of the fault.

An apparently anomalous feature of the zone is the lack of small earthquakes in recent years. Magnitude 4 events would have been completely reported since at least 1950, if not much earlier, magnitude 3 events since at least the early 1970's with the development of the Lamont-Doherty network in New York State. There have been very few such events in the last 15 years; consequently the data base for magnitude recurrence calculation is very sparse. There are only five Attica zone events that pass the completeness test (Figure 50 and Appendix A).


This is also an example of a zone that has experienced only one known significant earthquake in historic time, albeit only M 5.5, and it is therefore difficult to estimate the expected rate of such events. Consultations with G. Leblanc and P.W. Pomeroy (personal communications, 1981) have provided evidence that the settlement of western New York State was sufficiently dense by 1850 to have provided written accounts of Attica-sized earthquakes since that time. This date has been used (Table 1) to estimate rates of M 5.5 events for the magnitude recurrence relation (Figure 51). The maximum magnitude has been chosen as one-half magnitude unit larger, at 6.0.

3.24 Eastern Background (EBG) (Figures 52, 53)

The Eastern Background zone (Figure 52) has been drawn to encompass the entire region of eastern Canada that shows some evidence of minor historical, or recent low-level seismicity. As such, it extends beyond the more concentrated activity defined by the above zones and includes seismicity in regions of James Bay, northeast of Georgian Bay, western Lake Erie, the north shore of the Gulf of St. Lawrence and the northeastern Newfoundland Shelf. Defining the zone is primarily a recognition that low-level, but occasionally significant, seismicity can occur in regions surrounding the active zones; i.e., this region of eastern Canada should not be considered aseismic. The maximum magnitude is selected as 5.0, but the area is so large and the rates so low (Figure 53) that the zone makes a negligible contribution to the probabilistic ground motion in Figures 2 and 3. An earthquake in 1922 that may have had a magnitude near 5 has an epicentre on the northeastern Newfoundland shelf (see Figure 3 of Basham and Adams (1982)). This event does not pass the completeness test (Table 1) and its location is very uncertain.


3.25 Baffin Bay (BAB) (Figures 54, 55)

The largest earthquake known to have occurred in northern Canada was the magnitude 7.3 event in Baffin Bay in 1933 (Figure 54). This earthquake had aftershocks as large as M 6.5. Magnitude 6 events in Baffin Bay have since occurred in 1945, 1947 and 1957. It is difficult to define boundaries for the Baffin Bay zone on the basis of geological and geophysical evidence (Basham et al., 1977; Wetmiller and Forsyth, 1978, 1982; Reid and Falconer, 1982), but there seems to be a clear separation between the activity in the Bay and that on Baffin Island. Therefore, the zone boundaries shown in Figure 54 are



EARTH PHYSICS BRANCH
DIRECTION DE LA PHYSIQUE DU GLOBE EMR OTTAWA CANADA

Figure 52







Figure 54



EARTH PHYSICS BRANCH EMR OTTAWA CANADA
DIRECTION DE LA PHYSIQUE DU GLOBE OTTAWA CANADA

29/10/81 10.21.08.

Figure 55

arbitrary ones drawn to distinguish the significant seismicity of the Bay from surrounding events in the Baffin Island and Eastern Arctic Background zones discussed below.

Jackson et al. (1979) have found evidence for sea-floor spreading and an extinct spreading centre in the deep central region of the Bay. However, there is little or no seismic activity in this region; the seismicity is confined almost exclusively to the landward side of the 2000m bathymetric contour in the northwestern segment of the Bay that outlines the thick sedimentary sequence. This sedimentary sequence is also reflected by a broad positive free air gravity anomaly which suggests uncompensated loads may be acting on zones of weakness along the rifted margin (Wetmiller and Forsyth, 1982).

Stein et al. (1979) have found thrust mechanisms for the 1933 and a magnitude 5.6, 1976 earthquake in Baffin Bay and suggest that the stresses due to glacial unloading are sufficient to reactivate old faults parallel to the margin. They used synthetic seismogram calculations to suggest a 65-km focal depth for the 1933 earthquake, a surprisingly large value that must be considered poorly controlled because of the sparse seismic data available. Reid and Falconer (1982), however, employed the results of a microearthquake survey using ocean-bottom seismographs to make a speculative suggestion that current seismicity might be occurring on the deep 1933 thrust plane.

The magnitude recurrence relation for the Baffin Bay zone is reasonably well defined (Figure 55), but, as for the Laurentian Slope zone discussed above, the single large earthquake, when counted for the time period of complete reporting, produces too high a rate for that magnitude category. In this case a starting date of 1850 is imposed arbitrarily on the largest

magnitude category (Table 1). The assigned maximum magnitude is 7.5; thus, we suggest that the 1933 earthquake was near to the maximum size that can occur at any location in Canada away from active plate boundaries.

Because the zone is poorly defined and the seismicity dominated by one large earthquake, the implications of this model for seismic risk in Baffin Bay are similar to those discussed above for the Laurentian Slope zone.


3.26 Baffin Island (BAI) (Figures 56, 57)

Prior to 1960 only one earthquake is known to have occurred on Baffin Island, a moderate M5.5-6 event in 1935. With the development of the seismograph network in the north in the 1960's, in particular the station at Frobisher in 1963, the northeastern portion of the island was found to be highly active. This activity appears to be confined to the coastal region, and does not occur much further inland than the heads of the fjords, although there are scattered epicentres, perhaps mislocations, that extend to the centre of the island (see Figure 8 of Basham et al., 1977). The seismicity is concentrated in the regions of Buchan Gulf and Home Bay. There is a possible gap between these two concentrations, but, because of the short history and the nature of the seismicity, the gap is not recognized in drawing the crude rectangular boundary for the zone shown in Figure 56.

The seismicity tends to occur in swarms, with many events of similar magnitude, rather than as typical mainshock-aftershock sequences. All available evidence on focal depths suggests the earthquakes are shallow. Hashizume (1973) determined depths of 4-6 km for the 1970, M4.4 and 1972, M5.1 earthquakes; Liu and Kanamori (1980) a depth of 7 km for the 1963, M6.1



Figure 56



EARTH PHYSICS BRANCH EMR OTTAWA CANADA
DIRECTION DE LA PHYSIQUE DU GLOBE OTTAWA CANADA

29/10/81 10.21.08.

Figure 57

earthquake. Each of these earthquakes shows normal faulting, which Stein et al. (1979) attribute to reactivation of the basement faults by flexure caused by deglaciation. Basham et al. (1977) suggested that the centre of postglacial uplift over Foxe Basin is producing tilting of Baffin Island with a high differential uplift rate, or a hinge zone, in the region of seismicity along the northeastern coast.

The swarm-like nature of the Baffin Island seismicity results in a relatively large slope to the magnitude recurrence relation (Figure 57). The maximum magnitude selected is 7.0. This may be too large for shallow swarm seismicity but given the short earthquake history of the island it is considered a prudent choice.

3.27 Labrador Sea (LAB) (Figures 58, 59)

The known seismicity of the Labrador Sea includes six earthquakes in the M5.0 - 5.6 range (1934, 1952, 1956, 1958, 1962 and 1971, three of which pass the completeness test and are plotted in Figure 58), but none larger. There are reports of felt earthquakes from fishing villages along the Labrador coast as early as 1809 (Smith, 1962), and epicentres for these events have been assigned to the locations at which they were felt. However, there is no evidence from recent instrumental data that significant earthquakes are occurring onshore in this region. These older events likely occurred offshore and the zone defined here (Figure 58) is confined to the offshore region.

The Labrador Sea is a product of seafloor spreading and Srivastava (1978) has identified a central ridge and associated fracture zones from seismic and gravity profiles and linear magnetic anomalies. The central





Figure 58

ZONE: LABRADOR SEA (LAB)



EARTH PHYSICS BRANCH EMR OTTAWA CANADA
DIRECTION DE LA PHYSIQUE DU GLOBE OTTAWA CANADA

29/10/81 16.09.48.


Figure 58

Labrador Sea epicentres lie near, but southwest of, the ridge structures; i.e., the ridge appears to be the northeastern boundary of the activity rather than a locus. No earthquakes have been located between the ridge and Greenland (see Figure 2 of Basham and Adams, 1982). There is an apparently separate trend of earthquakes that follows the ocean-continent boundary northward from a fracture zone offshore from Hamilton Inlet, merging with the ridge trend near the northernmost seafloor ridge features. Thus, in general terms, the earthquakes must be associated with pre-existing faults near the inactive ridge and beneath the rifted continental margin. With further research, it may be possible to divide the source zone into two parts based on these trends. The best known earthquake is the M5.6, 1971 event (the southernmost epicentre in Figure 58) for which Hashizume (1977) determined a dip-slip mechanism at a depth of 16 km due to deviatoric compressive stresses normal to the margin.

The magnitude recurrence relation (Figure 59) is not well defined and the choice of maximum magnitude is a difficult one. A value of 6.5 has been chosen, but there is no evidence to suggest that the continental margin of the Labrador Sea cannot experience a magnitude 7 earthquake similar to the two that have occurred in historic time in the Baffin Bay and Laurentian Slope zones discussed above.

3.28 Eastern Arctic Background (EAB) (Figures 60, 61)

The region surrounding the Baffin Bay and Baffin Island zones (as defined above) has experienced low levels of both historic and recent seismicity. This includes the regions of the continental margin of Greenland, northern Davis Strait, the northern portion of Baffin Bay and Lancaster Sound




EARTH PHYSICS BRANCH
DIRECTION DE LA PHYSIQUE DU GLOBE EMR OTTAWA CANADA

Figure 60

ZONE: EASTERN ARCTIC BACKGROUND (EAB)

CUMULATIVE RATE (P.A.)



EARTH PHYSICS BRANCH EMR OTTAWA CANADA
DIRECTION DE LA PHYSIQUE DU GLOBE OTTAWA CANADA

29/10/81 16.09.48.

Figure 61

(Figure 60; see also Figure 2 of Basham et al., 1977). The Eastern Arctic Background zone, with rather arbitrary boundaries, is intended to account for this scattered seismicity.

Wetmiller and Forsyth (1982) have shown that Nares Strait, between Ellesmere Island and Greenland, is currently aseismic, but that a trend of epicentres from northern Baffin Bay appears to extend into Lancaster Sound. The Lancaster Sound events in their Figure 8 do not pass the completeness test for inclusion in Figure 60, although the northwestern boundary of the zone has been drawn to include this region. No earthquakes with magnitude greater than 4 have occurred in Davis Strait between the Eastern Arctic Background and the Labrador Sea zones since at least 1962 (Basham et al., 1977), and this region is considered to be aseismic in the present model (see Figure 5).

3.29 Gustaf-Lougheed Arch (GLA) (Figures 62, 63)

The Queen Elizabeth Islands seismicity is characterized by low to moderate magnitude, but often intense, earthquake swarms. The general cause of this seismicity is movement, suggested to be in response to the contemporary stress field, on unhealed faults that were formed or reactivated by Paleozoic and later orogenic phases (Basham et al., 1977; Forsyth et al., 1979; Wetmiller and Forsyth, 1982). However, because of the swarm-like nature of the seismicity, which has been observed to start abruptly in previously quiet areas, and the short observation period (about 20 years) it is highly unlikely that all potentially active regions of the islands have been identified.

A section through the Queen Elizabeth Islands that shows the highest levels of seismicity and for which there is some geological and geophysical





Figure 62

ZONE: GUSTAF-LOUGHEED ARCH (GLA)



EARTH PHYSICS BRANCH EMR OTTAWA CANADA
DIRECTION DE LA PHYSIQUE DU GLOBE OTTAWA CANADA

29/10/81 16.09.48.

Figure 63

evidence is defined here as the Gustaf-Lougheed Arch zone (Figure 62). The cluster of earthquakes in the Byam Martin Channel northeast of Melville Island occurred as intense swarms that started abruptly in 1972 (Basham et al., 1977). A similar but, to date, a less intense swarm occurred in Prince Gustaf Adolf Sea east of Borden Island in 1978. These latter events are included here, as an exception to the 1977 cut-off date used for a majority of the source zones, because they are important in defining the trend of the zone and are considered typical of the swarm activity to be expected along the zone in future. The zone is extended offshore to include the cluster of earthquakes on the continental slope north of Borden Island.

The name for this zone is taken from the Gustaf-Lougheed Arch, a structurally significant feature reflected in Bouguer gravity anomaly contours, that divides the western Sverdrup Basin into two separate sub-basins (Hea et al., 1979; Forsyth et al., 1979). The zone boundary in Figure 62 follows the outline of the arch from west of Elles Ringnes Island to southern Melville Island. Superimposed on the arch is a series of northeasterly trending minor magnetic highs reflecting mineralized faults or intrusive dykes.

The focal mechanisms for the four largest earthquakes in the Byam Martin Channel swarm (magnitudes 5.1 - 5.7) show deviatoric tension at depths from 9 km (just beneath the sediments of the Sverdrup Basin) to 31 km (Hasegawa, 1977), suggesting the fractures or dykes are loci of current seismic activity. A tensional regime along the Gustaf-Lougheed Arch suggests that current tectonic forces are similar to those in the Early Cretaceous responsible for the opening of the Arctic Ocean Basin (Sweeney et al., 1978).

There is little evidence to extend the Gustaf-Lougheed Arch, itself, offshore to the cluster of epicentres on the continental slope, but there is

other evidence of significant structures in the area. Submarine physiography indicates as much as 400 m of drowning over the continental shelf and slope offshore of Elles Ringnes Island, and submarine valleys on either side of the island trend toward the offshore cluster. This seismicity is on the seaward gradient of a large free air gravity anomaly, suggesting a region of stress adjustment to an uncompensated wedge of sediments (Basham et al., 1977).

There is no evidence on which to base a maximum magnitude (chosen as 6.5), but the choice does have an influence on the resulting probabilistic ground motion. The Gustaf Lougheed Arch zone produces a narrow zone of high acceleration (Figure 2), but a less pronounced zone of velocity (Figure 3).

3.30 Sverdrup (SVD) (Figures 64, 65)

With the Gustaf-Lougheed Arch zone (above) defined separately, the remainder of the Sverdrup Basin is seen as having a broad scattering of low level seismicity. It is characterized by both intense low-magnitude swarms such as that which occurred on Prince Patrick Island in 1965 (Smith et al., 1968), and single larger events with few detectable aftershocks such as the M 5.2 event on western Axel Heiberg Island in 1975. There are, however, numerous smaller earthquakes, that do not pass the completeness test used here, with epicentres in the Sverdrup Basin. The boundaries for the zone shown in Figure 64 follow, as closely as possible with long straight-line segments, the outer edge of the Franklinian province that surrounds the basin (see Figure 1 in Sweeney (1976)).

The small number of earthquakes that pass the completeness test produces difficulties in defining a magnitude recurrence relation. The curve in Figure 65 is defined by imposing the recurrence slope for the Gustaf-Lougheed Arch zone on the rates estimated from the small number of Sverdrup zone earthquakes.





Figure 64



EARTH PHYSICS BRANCH EMR OTTAWA CANADA
DIRECTION DE LA PHYSIQUE DU GLOBE OTTAWA CANADA

30/10/81 10.39.43.

Figure 65

3.31 Boothia-Ungava (BOU) (Figures 66, 67)

The seismicity of the Queen Elizabeth Islands is connected to the south by a concentration of epicentres in Barrow Strait south of Cornwallis Island, with a more diffuse trend through Somerset Island, down Boothia Peninsula and thence southeastward to a major cluster in the area of Wager Bay and Roes Welcome Sound. A less well-defined trend continues across the Ungava Peninsula and through Hudson Strait, connecting to the seismicity in the Labrador Sea. This seismicity is modelled here as one continuous narrow zone (Figure 66).

In general terms the seismicity in this arcuate band surrounds the southwestern half of the area of postglacial uplift centred over Foxe Basin, which led Basham et al., (1977) to speculate that the Baffin Island-Foxe Basin block is responding independently to postglacial uplift and may be decoupled from the rest of the shield to the southwest. Geological correlations are best at the north end where the seismicity shows a close relationship to the Boothia Uplift from Somerset and Prince of Wales Islands northward, meeting the Sverdrup Basin in the region of Grinnell Peninsula (Wetmiller and Forsyth, 1982). The Boothia Uplift, which has geologically demonstrated tectonic activity from the Paleozoic to the Cretaceous, continues to be active in present times. Seismological analysis shows that two earthquakes on the western edge of Southampton Island had focal depths of 17-21 km with thrust mechanisms due to northeast-southwest compression (Hashizume, 1974). The remaining seismicity through to the eastern end of Hudson Strait follows a broad deformational trend suggested by the Bell Arch and a series of horst-graben structures indicated in the bathymetry (Basham et al., 1977).





Figure 66

ZONE: BOOTHIA-UNGAVA (BOU)



EARTH PHYSICS BRANCH EMR OTTAWA CANADA
 DIRECTION DE LA PHYSIQUE DU GLOBE OTTAWA CANADA

29/10/81 16.09.48.

Figure 67

4. PROBABILISTIC SEISMIC GROUND MOTION ANALYSIS

The seismic risk calculations are made with an adaption of the McGuire (1976) computer program. This program has been modified for the Cyber system in the Energy, Mines and Resources Computer Science Centre and a program listing is included here as Appendix B. McGuire (1976) provides details of the functioning of the program and the various options available in its use. It is the purpose of this section to document special features of the program that are of concern in our application, some changes and additions that were made for this purpose, and the procedures used to make the calculations on a national-scale grid for purposes of contouring probabilistic peak acceleration and velocity (as shown in Figures 2 and 3).

4.1 Regionalization

There are two purposes for regionalizing the national-scale risk calculations: one is to accommodate the difference in strong seismic ground motion attenuation in western and eastern Canada; the other to avoid excessive calculations for a particular grid point by excluding source zones that will make no contribution to the resulting risk.

Hasegawa et al., (1981), in deriving separate strong motion attenuation relations for western and eastern Canada, defined a boundary between the two attenuation regions as following the eastern edge of Cordilleran tectonic province. This boundary is a distinctive physiographic feature at all locations along its length, except where it approaches the Beaufort Sea. The Beaufort Sea is included here in the western attenuation region, as it is viewed as being more analogous to the more recently tectonically active western region than to the less active eastern region. Thus, with reference

to Figure 5, it can be seen that all of the contiguous western source zones are in the western attenuation region, and all of the contiguous eastern source zones and the Southern Saskatchewan zone are in the eastern attenuation region. In one area only, the western Arctic islands region, are the two sets of source zones close enough to produce significant ground motion in the other region. This is handled as described below.

To accommodate the two attenuation regions and to avoid excessive calculations for source zones that make no contribution, the country has been divided into the eight regions described in Table 3. Each of the eight regions has a separate computational grid for calculations of contributions from the indicated source zones within the indicated latitude and longitude bounds. Some source zones are included for two or three regions because they make contributions outside of their own region.

In the "Northwest-Northeast" region the calculations are made assuming that the ground motion propagates from the western source zones with western attenuation into the eastern region, and from the eastern source zones with eastern attenuation into the western region. The computer program normally cumulates risk at fixed ground motion levels and then interpolates to produce ground motion at fixed risk levels. For this region there is a special version of the program in which the two sets of risk cumulations are done separately, and then added together prior to the interpolation to produce ground motion at fixed risk levels.

4.2 Integration over Source Zones

Source zones and site locations are defined by geographical coordinates given in degrees of latitude and longitude. As the McGuire program was designed as a planar version using Cartesian coordinates, all of the

Table 3

Regionalization for Risk Calculationsa) Region of eastern attenuation

1. "East": 41.0 to 50.0°N, 49.0 to 86.0°W
Source zones: CHV, WQU, LSL, NAP, ATT, LSP and EBG
2. "East-North": 50.0 to 60.0°N, 50.0 to 90.0°W
Source zones: CHV, WQU, LSL, NAP, LSP, EBG, LAB, GLA, SVD, BOU, BAI, BAB and EAB
3. "North-East": 60.0 to 85.0°N, 50.0 to 105.0°W
Source zones: GLA, SVD, BOU, BAI, BAB, LAB and EAB
4. "Central": 46.0 to 51.0°N, 100.0 to 110.0°W
Source zones: SAS, SBC, and FHL

b) Region of western attenuation

5. "West": 47.0 to 53.0°N, 109.0 to 142.0°W
Source zones: CAS, PGT, CSM, NVI, JFE, QCF, SPT, SBC, NBC and FHL
6. "West-North": 53.0 to 60.0°N, 110.0 to 145.0°W
Source zones: CAS, PGT, CSM, NVI, JFE, SBC, FHL, QCF, SPT, NBC, FWY, DSK, RIC, MKZ, BFT, ALC and ALI
7. "North-West": 60.0 to 68.0°N, 110.0 to 145.0°W
Source zones: FWY, DSK, RIC, MKZ, BFT, QCF, SPT, NBC, ALC and ALI

c) Region requiring eastern and western attenuation

8. "Northwest-Northeast": 68.0 to 82.0°N, 91.0° to 145.0°W
Source zones with western attenuation: FWY, DSK, RIC, MKZ, BFT, QCF, SPT, NBC, ALC and ALI

Source zones with eastern attenuation: GLA, SVD, BOU, BAI, BAB, LAB and EAB.

geographic coordinates are transformed into eastings and northings in kilometres using a Lambert Conformal projection, with each region given its own central meridian. Any distortion in the calculated distances is well below the accuracy of the distances required in estimating ground motion.

In integrating the contributions to the risk at a site from a source zone, the source zone is divided into finite arc segments, the radii of which pivot on the site. The size of the arc segments is a function of the gross source dimensions and the program parameter NSTEP. As NSTEP is increased the area covered by the arc segments tends to the correct area of the source zone. A value of NSTEP = 10 is found to be adequate in most cases, as increasing it does not significantly change the calculated values. However, the computation time does increase with NSTEP so NSTEP is kept as small as possible, the limitation being the area error that will be tolerated. For some site-source combinations, a value of NSTEP = 10 leads to significant area errors. If the area error is greater than ERRBND percent (20 is used in current applications) then NSTEP is automatically doubled and the calculation repeated until either the error is less than ERRBND or NSTEP reaches the assigned maximum value of NSTEPMX. If NSTEP reaches NSTEPMX, the flag LERR is set for the calculation. After contributions from all source zones are integrated, the ratio of the flagged to unflagged risk cumulations is computed. If this ratio is less than RKRATO (0.05 is used in current applications), the result is accepted; otherwise the total risk for the site is flagged. When contouring risk computed on a grid (see section 4.6), flagged grid points are omitted.

The program parameter RZ2 defines the closest distance from site to source zone beyond which the earthquakes are considered lumped at the centre

of the source zone. In the original program RZ2 was fixed at 300 km with no provision for redefinition. To allow for various source zone geometries for which $RZ2 = 300$ is not the most appropriate, a change has been made to read in this parameter with the other source zone parameters. In the current application $RZ2 = 600$ has been used for the following source zones: FWY, ALC, ALI, QCF, SPT, LAB, GLA and BOU.

4.3 Treatment of Other Errors

In subroutines INSIDE and OUTSIDE some types of errors can occur which are a function of the site-source geometry and result in no risk being computed. For example, the distance to the nearest zone boundary may become zero. Again NSTEP is increased as described above to try to eliminate the problem. If the error persists LERR2 is set "true" and the result is flagged.

In all cases a message is printed when these errors occur giving the site location and the source zone in which the error is encountered. If an output is desired in order to see what values are being calculated, the omission of erroneous results can be cancelled by setting INCLUD and/or INCLUD2. Re-definition of source zone geometries in the area of the site will usually be necessary if a valid estimate is required in these cases.

In the current calculations for Figures 2 and 3, only three grid points, out of more than 6000, had to be omitted. These were at 51.0°N , 122.0°W , at the junction of the CSM, SBC and CAS zones in southern British Columbia; at 61.0°N , 145.0°W , at the junction of the FWY, DSK, ALI and ALC zones in Alaska; and at 67.5°N , 67.0°W , at the common corner of the BAI and EAB zones on Baffin Island (see Figure 5). Five other grid points had area errors greater than 20 percent for $\text{NSTEP}=10$, but these were reduced to less than 20 percent by the automatic recalculation with NSTEP increased.

4.4 Limiting Ground Motion from Large Magnitudes

As noted in Section 2.3, the expressions for strong seismic ground motion attenuation derived by Hasegawa et al. (1981) did not limit ground motion contributions at large magnitudes. For current computations this has been implemented in the program (see subroutine RISK1 in Appendix B) by modifying the magnitude recurrence relation so that all expected events greater than M7.5 are compressed into a Delta function (a spike) at M7.5.

4.5 Statistical Scatter on Attenuation

The program provides for the inclusion of a standard deviation on the logarithm of the ground motion parameter whose mean value is defined by the attenuation relation, and a normal distribution is assumed. In the current application this variable (SIG) is set to 0.7, the natural logarithm of 2; i.e., the standard deviation on both peak acceleration and velocity are assumed to be a factor of 2.

4.6 Computation Grid and Contouring

The ground motion values for Figures 2 and 3 were computed using the grid of points listed in Table 4. This grid has a maximum spacing of 57 km in latitude and 90 km in longitude, with progressively smaller longitude spacing to the north. The computations were stored on a computer file containing latitude, longitude, risk values and their corresponding acceleration and velocity values.

The contouring of the data was done in five regions, east, northeast, west, northwest and central. For each region the data from the risk program were first combined through a program that projects latitude and longitude

Table 4

Computation Grid used for Contouring

<u>Map Sheet</u>	<u>Lat. Boundaries</u>	<u>Step</u>	<u>Long. Boundaries</u>	<u>Step</u>
1	41.0 - 49.5	0.5	86.0 - 50.0	1.0
2	50.0 - 59.5	0.5	90.0 - 50.0	1.0
3	60.0 - 62.5	0.5	95.0 - 50.0	1.0
	63.0 - 65.5	0.5	100.0 - 50.0	1.0
	66.0 - 67.5	0.5	105.0 - 50.0	1.0
	68.0 - 79.5	0.5	90.0 - 50.0	1.0
	80.0 - 85.0	1.0	90.0 - 50.0	2.0
4	68.0 - 79.5	0.5	145.0 - 91.0	1.0
	80.0 - 85.0	1.0	145.0 - 90.0	2.0
5	60.0 - 67.5	0.5	145.0 - 110.0	1.0
6	53.0 - 59.5	0.5	145.0 - 110.0	1.0
7	47.0 - 52.5	0.5	142.0 - 110.0	1.0
8	42.0 - 46.5	0.5	137.0 - 110.0	1.0
9	46.0 - 51.0	0.5	110.0 - 101.0	1.0
10	66.0 - 67.5	0.5	109.0 - 106.0	1.0
	63.0 - 65.5	0.5	109.0 - 101.0	1.0
	60.0 - 62.5	0.5	109.0 - 96.0	1.0

into eastings and northings (x and y) on a Lambert projection at a fixed scale. The input data were selected so as to extend beyond the area to be contoured in order to obtain continuity in the contours between regions and to avoid edge effects. The output from this program was formatted so as to be compatible with the program GPCP described below.

The contouring was done with the "General Purpose Contouring Program" (GPCP), a product of the Calcomp Company, which resides on the EM&R Computer Science Centre Cyber system. By using the x and y dimensions obtained from the projection program, the contours are properly scaled to the scale selected for the map area. The GPCP program requires the x-y grid to be specified so that each cell contains no more than one data point. Cell size can be calculated knowing the latitude-longitude grid, as specified in Table 4, and the x and y dimensions as provided by the projection program.

The method of contouring is described in detail by the GPCP users guide prepared by the Computer Science Centre. Briefly, the method is as follows. Using data supplied on the "SIZX" input card, a uniform x-y grid is established, the data at each grid point being approximated by a function defined by the nearest "n" data points. "n" can be defined on the "CNTL" card; the default value of n=8 was used. To generate smooth contours, each grid cell is divided into a finer sub-grid using a third-order interpolation, and the contour lines are drawn as short straight-line segments between the sub-grid points. The default value of 5 was used to divide both the x and y sides of each grid cell.

The final contour maps were drafted from the five partially-overlapping region maps. Some "chatter" in the contours occurs due to the finite grid spacing of the calculations (Table 4), which is smoothed by hand during

drafting. This "chatter" could be removed by a denser original grid, but the extra computations are not considered justified for the resolution required in the final maps.

4.7 Site-Specific Risk Calculations

The earthquake source zones as defined in Section 3 above and the computer program listed in Appendix B will be maintained by the Earth Physics Branch on the Computer Science Centre Cyber system. This package of input data and computation method will remain intact for some years in order that computations can be made at any time, equivalent to those used for the contour maps recommended as new zoning maps for the National Building Code.

Using this package, the Earth Physics Branch will undertake site-specific risk calculations on request for a nominal fee. A sample of the output information for a site-specific calculation is given in Table 5. The request will specify a site with geographical coordinates. The output will be peak acceleration and velocity for the four risk levels indicated. The risk level of 0.002105 per annum is equivalent to a probability of 10 percent exceedence in 50 years (e.g., Figures 2 and 3) that has been recommended for the National Building Code zoning maps.

ACKNOWLEDGEMENTS

We benefited greatly from numerous discussions with our Earth Physics Branch colleagues during the preparation of this report, in particular, W.G. Milne (now with Woodward Clyde Consultants), R.D. Hyndman, J. Adams, A.E. Stevens, R.B. Horner, and G.C. Rogers. J.A. Drysdale provided valuable assistance with the early preparation of earthquake lists and source zone maps. R.B. Horner performed the recomputations of magnitudes and A.E. Stevens reviewed a number of historical earthquakes, the results of which were employed in preparing Appendix C.

ENERGY, MINES AND
RESOURCES CANADA
EARTH PHYSICS BRANCH

ENERGIE, MINES ET
RESSOURCES CANADA
DIRECTION DE LA PHYSIQUE DU GLOBE

SEISMIC RISK CALCULATION *

CALCUL DE RISQUE SEISMIQUE *

REQUESTED BY

SEISMIC HAZARDS AND APPLICATIONS EPB/EMR

DEMANDE PAR

FOR SITE

OTTAWA, CANADA

POUR SITE

LOCATED AT

45.39 NORTH/NORD

75.72 WEST/OUEST

LOCATION

PROBABILITY OF EXCEEDENCE
PER ANNUM

.01

.005

.002105

.001

PROBABILITE DE
DEPASSEMENT PAR ANNEE

PEAK HORIZONTAL
ACCELERATION (%G)

8.4

12.3

20.0

28.3

ACCELERATION HORIZONTALE
MAXIMALE (%G)

PEAK HORIZONTAL
VELOCITY (CM/SEC)

3.1

5.4

9.8

15.2

VITESSE HORIZONTALE
MAXIMALE (CM/SEC)

* REFERENCE

NEW PROBABILISTIC STRONG SEISMIC GROUND
MOTION MAPS OF CANADA: A COMPILATION OF EARTHQUAKE
SOURCE ZONES, METHODS AND RESULTS
P.W. BASHAM, D.H. WEICHERT, F.M. ANGLIN, AND M.J. BERRY
EARTH PHYSICS BRANCH OPEN FILE NUMBER 82-
OTTAWA, CANADA 1982

09/12/82 12.48.13.

REFERENCES

- Aggarwal, Y.P., and L.R. Sykes (1978). Earthquakes, faults and nuclear power plants in southern New York and northern New Jersey. *Science*, 200, 425-429.
- Algermissen, S.T. and D.M. Perkins (1976). A probabilistic estimate of maximum acceleration in rock in the contiguous United States. U.S. Geol. Surv. Open File Rept. 76-416, 45 pp.
- Anglin, F.M., and G. Buchbinder (1981). Microseismicity in the mid-St. Lawrence Valley Charlevoix zone, Quebec. *Bull. Seism. Soc. Am.* 71, 1553-1560.
- Associate Committee on the National Building Code (1980). National Building Code of Canada 1980, NRCC No. 17303; The Supplement to the National Building Code of Canada 1980, NRCC No. 17724, National Research Council, Ottawa.
- Basham, P.W., and K. Whitham (1966). Microseismic noise on Canadian seismograph records in 1962 and station capabilities. *Pub. Dom. Obs.*, Ottawa, 32, 123-135.
- Basham, P.W. and D.H. Weichert (1979). Seismic risk mapping in Canada. *Proc. Third Can. Conf. Earthquake Eng.* 1, 23-48, Montreal.
- Basham, P.W. and J. Adams (1982). Earthquake hazards to offshore development on the eastern Canadian continental shelves. *Proc. Second Can. Conf. Marine Geotech. Eng.*, XX-XX, Dartmouth.
- Basham, P.W., D.A. Forsyth and R.J. Wetmiller (1977). The seismicity of northern Canada. *Can. J. Earth Sci.* 14, 1646-1667.
- Basham, P.W., D.H. Weichert and M.J. Berry (1979). Regional assessment of seismic risk in eastern Canada. *Bull. Seism. Soc. Am.* 69, 1567-1602.
- Basham, P.W., P. Morel-à-l'Huissier and F.M. Anglin (1982). Earthquake risk at Gros Cacouna, Quebec, and Melford Point, Nova Scotia. *Earth Physics Branch Open File Rept.* 82-2, 116 pp.
- Basham, P.W., D.H. Weichert, F.M. Anglin and M.J. Berry (1983). New probabilistic seismic ground motion maps of Canada. *Bull. Seism. Soc. Am.* (submitted for publication).
- Berry, M.J., F.M. Anglin and P.W. Basham (1982). Earthquake risk in eastern Canada. *Proc. Internat. Symposium on Continental Seismicity and Earthquake Prediction, Beijing, China* (in press).
- Bird, D.N. (1981). Time-term analysis using linear programming and its application to refraction data from the Queen Charlotte Islands. M.Sc. Thesis, University of British Columbia, Vancouver, 79 p.

- Campbell, K.W. (1981). Near-source attenuation of peak horizontal acceleration. *Bull. Seism. Soc. Am.* 71, 2039-2070.
- Coombs, H.A., W.G. Milne, O.W. Nuttli, and D.B. Slemmons (1976). Report of the review panel on the December 14, 1872 earthquake. Pacific Northwest Utilities, Special Rept.
- Cornell, C.A. (1968). Engineering seismic risk analysis. *Bull. Seism. Soc. Am.* 58, 1583-1606.
- Crosson, R.S. (1982). Review of seismicity in the Puget Sound region from 1970 through 1978. U.S. Geol. Surv. Open File Report .
- Currie, R.G., L.E. Stevens, D.L. Tiffin and R.P. Riddihough (1980). Marine geophysical survey of the Queen Charlotte Islands (abstract) EOS, 61, p. 71.
- Davis, E.E., and R.P. Riddihough (1982). The Winona Basin: structure and tectonics. *Can. J. Earth Sci.* 19, 767-788.
- Doxsee, W.W. (1948). The Grand Banks earthquake of November 18, 1929. *Pub. Dom. Obs.* 7, 323-335.
- Fletcher, J.B., and L.R. Sykes (1977). Earthquakes related to hydraulic mining and natural seismic activity in western New York State. *J. Geophys. Res.* 82, 3767-3780.
- Forsyth, D.A. (1981). Characteristics of the western Quebec seismic zone. *Can. J. Earth Sci.* 18, 103-119.
- Forsyth, D.A., P. Morel, H.S. Hasegawa, R.J. Wetmiller, J. Adams, A.K. Goodacre, R.L. Coles, D. Nagy, J. Harris and P.W. Basham (1982). Comparative study of the geophysical and geological information in the Timiskaming-Kapuskasing Area. (in preparation).
- Forsyth, D.A. J.A. Mair and I. Fraser (1979). Crustal structure of the central Sverdrup Basin. *Can. J. Earth Sci.* 16, 1581-1598.
- Gendzwill, D.J., R.B. Horner and H.S. Hasegawa (1982). Induced earthquakes at a potash mine near Saskatoon, Canada. *Can. J. Earth Sci.* 19, 466-475.
- Goodacre, A.K., and H.S. Hasegawa (1980). Gravitationally induced stresses at structural boundaries. *Can. J. Earth Sci.* 17, 1286-1291.
- Gutenberg, B., and C.F. Richter (1954). Seismicity of the Earth and Associated Phenomena, 2nd. Ed., Princeton University Press, 310 pp.
- Hasegawa, H.S. (1977). Focal parameters of four Sverdrup Basin, Arctic Canada, earthquakes in November and December of 1972. *Can. J. Earth Sci.*, 11, 2481-2494.

- Hasegawa, H.S., and R.J. Wetmiller (1980). The Charlevoix earthquake of 19 August 1979 and its seismo-tectonic environment. *Earthquake Notes*, 51, 23-37.
- Hasegawa, H.S., and J. Adams (1981). Crustal stresses and seismotectonics in eastern Canada. *Earth Physics Branch Open File Rept.* 81-12, 62 pp.
- Hasegawa, H.S., C.W. Chou and P.W. Basham (1979). Seismotectonics of the Beaufort Sea. *Can. J. Earth Sci.* 16, 816-830.
- Hasegawa, H.S., P.W. Basham and M.J. Berry (1981). Attenuation relations for strong seismic ground motion in Canada. *Bull. Seism. Soc. Am.* 71, 1943-1962.
- Hasegawa, H.S., P.W. Basham and M.J. Berry (1982). Reply to Atkinson's "Discussion of 'Attenuation relations seismic ground motion in Canada'." *Bull. Seism. Soc. Am.* 72, 1771-1774.
- Hashizume, M. (1973). Two earthquakes on Baffin Island and their tectonic implications. *J. Geophys. Res.*, 78, 6069-6081.
- Hashizume, M. (1974). Surface wave study of earthquakes near northwestern Hudson Bay, Canada. *J. Geophys. Res.*, 79, 5458-5468.
- Hashizume, M. (1977). Surface-wave study of the Labrador Sea earthquake 1971 December. *Geophys. J.R. Astr. Soc.*, 51, 149-168.
- Hea, J.P., J. Arcuri, G.R. Campbell, I. Fraser, M.O. Fuglem, J.J. O'Bertos, D.R. Smith and M. Zayat (1979). Post Ellesmerian basins of Arctic Canada: their depocentres, rates of sedimentation and petroleum potential. In: *Facts and Principles of World Petroleum Occurrence* (A.D. Miall, Ed.). *Can. Soc. Petrol. Geol.*, Memoir 6.
- Heidebrecht, A.C., P.W. Basham, J.H. Rainer and M.J. Berry (1983). Engineering applications of new probabilistic seismic ground motion maps of Canada. *Can. J. Civil Eng.* (submitted for publication).
- Hodgson, J.H. (1956). A seismic probability map for Canada. *Can. Underwriter* 23, 7.
- Horner, R.B. (1983). Seismicity in the St. Elias region of northwestern Canada and southeastern Alaska. *Bull. Seism. Soc. Am.* (submitted).
- Horner, R.B., and H.S. Hasegawa (1978). The seismotectonics of southern Saskatchewan. *Can. J. Earth Sci.* 15, 1341-1355.
- Horner, R.B., A.E. Stevens and H.S. Hasegawa (1973). The Bengough, Saskatchewan, earthquake of July 26, 1972. *Can. J. Earth Sci.* 10, 1805-1821.
- Hyndman, R.D., and G.C. Rogers (1981). Seismicity surveys with ocean bottom seismographs off western Canada. *J. Geophys. Res.* 86, 3867-3880.

- Hyndman, R.D., and R.M. Ellis (1981). Queen Charlotte fault zone: microearthquakes from a temporary array of land stations and ocean bottom seismographs. *Can. J. Earth Sci.* 18, 776-788.
- Hyndman, R.D., and D.H. Weichert (1983). Seismicity and rates of relative motion on the plate boundaries of western North America. *Geophys. J.R.A.S.* (in press).
- Hyndman, R.D., G.C. Rogers, M.N. Bone, C.R.B. Lister, U.S. Wade, D.L. Barrett, E.E. Davis, T. Lewis, T. Lynch and D. Seemann (1978). Geophysical measurements in the region of Explorer Ridge off western Canada. *Can. J. Earth Sci.* 15, 1508-1525.
- Hyndman, R.D., R.P. Riddihough and R. Herzer (1979). The Nootka fault zone -a new plate boundary off western Canada. *Geophys. J. R.A.S.* 58, 667-683.
- Hyndman, R.D., T.J. Lewis, J.A. Wright, M. Burgess, D.S. Chapman and M. Yamano (1982). Queen Charlotte Fault Zone: heat flow measurements. *Can. J. Earth Sci.* 19, 1657-1669.
- Jackson, H.R., C.E. Keen, R.K.H. Falconer and K.P. Appleton (1979). New geophysical evidence for sea-floor spreading in Baffin Bay. *Can. J. Earth Sci.*, 16, 2122-2135.
- Joyner, W.B., and D.M. Boore (1981). Peak horizontal acceleration and velocity from strong-motion records including records from the 1979 Imperial Valley, California, earthquake. *Bull. Seism. Soc. Am.* 71, 2011-2038.
- Kanamori, H., and D.L. Anderson (1975). Theoretical basis of some empirical relations in seismology. *Bull. Seism. Soc. Am.* 65, 1073-1095.
- King, L.H. (1979). Aspects of regional surficial geology related to site investigation requirements - eastern Canadian shelf. *In*: Andus, D.A., (Ed.), Offshore Site Investigation, Graham and Trotman Ltd., London, 37-59.
- Klotz, O.J. (1915). Earthquake of February 10, 1914. *Pub. Dom. Obs. Ottawa*, 3, 3-14.
- Lahr, J.C., and G. Plafker (1980). Holocene Pacific-North America plate interaction in southern Alaska: implications from the Yakataga seismic gap. *Geology* 8, 483-486.
- Lahr, J.C., C.D. Stevens, H.S. Hasegawa and J. Boatwright (1980). Alaska seismic gap only partially filled by 28 February 1979 earthquake. *Science* 207, 1351-1353.
- Leblanc, G. (1981). A closer look at the September 16, 1732, Montreal earthquake. *Can. J. Earth Sci.* 18, 539-550.

- Leblanc, G., and R.J. Wetmiller (1974). An evaluation of seismological data available for the Yukon Territory and Mackenzie Valley. *Can. J. Earth Sci.* 11, 1435-1454.
- Leblanc, G., and G. Buchbinder (1977). Second microearthquake survey of the St. Lawrence Valley near La Malbaie, Quebec. *Can. J. Earth Sci.* 14, 2778-2789.
- Leblanc, G., and F.M. Anglin (1978). Induced seismicity at the Manic 3 reservoir, Quebec. *Bull. Seism. Soc. Am.*, 68, 1469-1485.
- Liu, H.-L., and H. Kanamori (1980). Determination of source parameters of mid-plate earthquakes from the waveforms of body waves. *Bull. Seism. Soc. Am.*, 70, 1989-2004.
- Malone, S.D., and S.-S. Bor (1979). Attenuation patterns in the Pacific Northwest based on intensity data and the location of the 1872 North Cascades earthquake. *Bull. Seism. Soc. Am.* 69, 531-546.
- McGuire, R.K. (1976). FORTRAN computer program for seismic risk analysis. U.S. Geol. Surv. Open File Rept. 76-67, 90 pp.
- McLellen, J.D., H.S. Hasegawa, J.-C. Roegiers and A.M. Jessop (1983). A hydraulic fracturing experiment at the University of Regina campus; geothermal and seismotectonic implications. *Can. Geotech. J.* (submitted for publication).
- Milne, W.G., and A.G. Davenport (1969). Distribution of earthquake risk in Canada. *Bull. Seism. Soc. Am.* 59, 729-754.
- Milne, W.G., G.C. Rogers, R.P. Riddihough, R.D. Hyndman and G.A. McMechan (1978). Seismicity of western Canada. *Can. J. Earth Sci.* 15, 1170-1193.
- Minster, J.B., and T.H. Jordan (1978). Present day plate motions. *J. Geophys. Res.* 83, 5331-5354.
- Norris, D.K. (1972). Geological map of the western part of the excursion area. 24th Int. Geol. Congr. Field Excursion A14, Guidebook, by A.C. Lenz and A.E.H. Pedder, 43 p.
- Perez, O.J., and K.H. Jacob (1980). Tectonic model and seismic potential of the eastern Gulf of Alaska and Yakataga seismic gap. *J. Geophys. Res.* 85, 7132-7150.
- Reid, I., and R.K.H. Falconer (1982). A seismicity study in northern Baffin Bay. *Can. J. Earth Sci.*, 19, 1518-1531.
- Riddihough, R.P. (1977). A model for recent plate interactions off Canada's west coast. *Can. J. Earth Sci.* 14, 384-396.
- Riddihough, R.P., R.G. Currie and R.D. Hyndman (1980). The Dellwood Knolls and their role in triple junction tectonics off northern Vancouver Island. *Can. J. Earth Sci.* 17, 577-593.

- Rogers, G.C. (1981). McNaughton Lake seismicity - more evidence for an Anahim hotspot? *Can. J. Earth Sci.* 18, 826-828.
- Rogers, G.C. (1982). Some comments on the seismicity of the northern Puget Sound-southern Vancouver Island region. In: *Earthquake Hazards of the Puget Sound Region, Washington State* (J.C. Yount, ed.) U.S. Geol. Surv. Open File Rept. (in press).
- Rogers, G.C. (1983). Revised seismicity and revised fault plane solutions for the Queen Charlotte Islands region and their tectonic implications. *Earth Physics Branch Open File Rept.* (in preparation).
- Rogers, G.C., and H.S. Hasegawa (1978). A second look at the British Columbia earthquake of June 23, 1946. *Bull. Seism. Soc. Am.* 68, 653-675.
- Rogers, G.C., and R.M. Ellis (1979). The eastern British Columbia earthquake of February 4, 1918. *Can. J. Earth Sci.* 15, 1484-1493.
- Rogers, G.C., R.M. Ellis and H.S. Hasegawa (1980). The McNaughton Lake earthquake of May 14, 1978. *Bull. Seism. Soc. Am.* 70, 1771-1786.
- Rondot, J. (1979). Reconnaissance Géologiques dans Charlevoix-Saguenay. *Ministère des Richesses naturelles du Québec*, DPV-682, 44 pp.
- Singh, S.K., E. Bozan and L. Esteva (1980). Expected earthquake magnitude from a fault. *Bull. Seism. Soc. Am.* 70, 903-914.
- Slawson, W.F., and J.C. Savage (1979). Geodetic deformation associated with the 1946 Vancouver Island, Canada, earthquake. *Bull. Seism. Soc. Am.* 69, 1487-1496.
- Smith, R.B. and Sbar, M.L. (1974). Contemporary tectonics and seismicity of the western United States with emphasis on the Intermountain seismic belt. *Bull. Geol. Soc. Am.* 85, 1205-1218.
- Smith, W.E.T. (1962). Earthquakes of eastern Canada and adjacent areas: 1534-1927. *Pub. Dom. Obs.* 26, 271-301.
- Smith, W.E.T. (1966). Earthquakes of eastern Canada and adjacent areas: 1928-1959. *Pub. Dom. Obs.* 32, 87-121.
- Smith, W.E.T., K. Whitham and W.T. Piché (1968). A microearthquake swarm in 1965 near Mould Bay, N.W.T., Canada. *Bull. Seism. Soc. Am.*, 58, 1991-2011.
- Srivastava, S.P. (1978). Evolution of the Labrador Sea and its bearing on the early evolution of the North Atlantic. *Geophys. J.R. Astro. Soc.*, 52, 313-357.
- Stein, S., N.H. Sleep, R.J. Geller, S.-C. Wang and G.C. Koeger (1979). Earthquakes along the passive margin of eastern Canada. *Geophys. Res. Letters*, 6, 537-540.

- Stevens, A.E. (1980). Re-examination of some larger La Malbaie, Quebec, earthquakes (1924-1978). *Bull. Seism. Soc. Am.* 70, 529-557.
- Stevens, A.E. (Editor) (1982). Preliminary report of the Miramichi, New Brunswick, Canada earthquake sequence of 1982. Earth Physics Branch Open File 82-24, 94 pp.
- Street, R.L., and F.T. Turcotte (1977). A study of northeastern North American spectral moments, magnitudes and intensities. *Bull. Seism. Soc. Am.*, 67, 599-614.
- Sweeney, J.F., (1976). Evolution of the Sverdrup Basin, Arctic Canada. *Tectonophysics*, 36, 181-196.
- Sweeney, J.F., E. Irving and J.W. Geuer (1978). Evolution of the Arctic Basin. *In* Arctic Geophysical Review (J.F. Sweeney, Ed.), Pub. Earth Physics Branch, 45, 91-100.
- Sykes, L.R. (1971). Aftershock zones of great earthquakes, seismicity gaps, and earthquake prediction for Alaska and the Aleutians. *J. Geophys. Res.* 76, 8021-8041.
- Thenhaus, P.C., J.I. Ziony, W.H. Diment, M.G. Hopper, D.M. Perkins, S.L. Hanson and S.T. Algermissen (1979). Probabilistic estimates of maximum seismic acceleration in rock in Alaska and the adjacent outer continental shelf. U.S. Geol. Surv. Interagency Rept. to the Bureau of Land Management.
- Uzumeri, S.M., S. Otani and M.P. Collins (1978). An overview of Canadian code requirements for earthquake resistant concrete buildings. *Can. J. Civil Eng.* 5, 427-441.
- von Heune, R., G.G. Shor and J. Wageman (1979). Continental margins of the eastern Gulf of Alaska and boundaries of tectonic plates. *In*: Geological and Geophysical Investigations of Continental Margins (J.S. Watkins, L. Montadert and P.W. Dickerson, Eds.), Am. Assoc. Petrol. Geol., Tulsa, Okla., 273-290.
- Weichert, D.H. (1980). Estimation of the earthquake recurrence parameters for unequal observation periods for different magnitudes. *Bull. Seism. Soc. Am.* 70, 1337-1346.
- Weichert, D.H., and W.G. Milne (1979). On Canadian methodologies of probabilistic seismic risk estimation. *Bull. Seism. Soc. Am.* 69, 1549-1566.
- Weichert, D.H., and R.D. Hyndman (1982a). A comparison of the rate of seismic activity and several estimates of deformation in the Puget Sound area. *In*: Earthquake Hazards of the Puget Sound Region, Washington State (J.C. Yount, Ed.). U.S. Geol. Surv. Open File Rept. (in press)

- Weichert, D.H., and R.D. Hyndman (1982b). Microzonation of the Queen Charlotte Islands and adjacent areas. Proc. Third Internat. Earthquake Microzonation. Conf., 3, 1489-1500.
- Wetmiller, R.J. (1975). The Quebec-Maine border earthquake, 15 June 1973. Can. J. Earth Sci. 12, 1917-1928.
- Wetmiller, R.J., and D.A. Forsyth (1978). Seismicity of the Arctic, 1908-1975. In Arctic Geophysical Review (J.F. Sweeney, Ed.), Pub. Earth Physics Branch, 45, 15-24.
- Wetmiller, R.J., and D.A. Forsyth (1982). Review of seismicity and other geophysical data near Nares Strait. In Nares Strait and the drift of Greenland: a conflict in plate tectonics (P.R. Dawes and J.W. Kerr, Eds.) Meddr. Gronland Geosci., 8, (in press).
- Wetmiller, R.J., J. Adams, A.E. Stevens, F.M. Anglin, H.S. Hasegawa and J. Bérubé (1982). Aftershock sequences for the New Brunswick earthquakes of January 9th and 11th, March 31st and June 16th, 1982 (Abstract). Earthquake Notes, (in press).
- Whitham, K., W.G. Milne and W.E.T. Smith (1970). The new seismic zoning map for Canada, 1970 edition. Can. Underwriter, 15, 6.
- Yang, J.-P., and Y.P. Aggarwal (1981). Seismotectonics of northeastern United States and adjacent Canada. J. Geophys. Res. 86, 4981-4998.
- Yorath, C.J., and R.L. Chase (1981). Tectonic history of allochthonous terrains, northern Canadian Pacific continental margin. Can. J. Earth Sci., 18, 1717-1739.
- Yorath, C.J., and R.D. Hyndman (1983). Subsidence and thermal history of Queen Charlotte Basin. Can. J. Earth Sci., 19 (in press).

APPENDIX A

Earthquakes that pass the completeness test of Table 1, that are employed to estimate magnitude recurrence relations, and that are plotted on each of the source zone maps, are listed on the following pages.

PUGET SOUND (PGT)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1909	01	11	23	49		48.7	122.8	6.0
1939	11	13	7	45	53	47.50	122.40	5.7
1943	11	29	01	43		48.40	122.90	5.0
1946	2	15	3	14	50	47.30	122.90	6.3
1949	4	13	19	55	36	47.20	122.60	7.1
1954	5	15	13	2	32	47.40	122.50	5.0
1957	01	26	01	16	07	48.29	122.60	5.0
1960	9	10	15	6	34	47.50	122.70	4.9
1962	12	31	20	49	35	47.00	122.00	5.0
1963	1	24	21	43	0	47.40	122.10	5.0
1964	10	14	6	33	0	47.70	122.10	4.6
1965	4	29	15	28	44	47.40	122.30	6.5
1965	10	23	16	28	3	47.50	122.40	4.8
1967	3	7	3	51	8	47.70	122.70	4.1
1967	05	25	23	22	35	48.20	122.81	4.3
1968	6	19	5	51	43	47.20	122.50	4.0
1968	9	6	12	16	30	47.95	122.80	3.9
1969	2	14	8	33	36	48.94	123.07	4.3
1970	5	18	5	29	54	48.60	122.70	4.0
1971	1	25	21	37	53	48.70	123.00	3.8
1971	12	28	7	50	0	47.60	122.20	4.0
1972	11	9	4	19	17	48.24	123.55	3.9
1974	5	16	13	4	36	48.14	122.92	3.8
1975	4	23	1	3	43	47.11	122.63	3.8
1976	5	16	8	35	15	48.80	123.34	5.4
1976	9	2	13	36	11	48.19	122.72	4.5
1976	9	8	8	21	2	47.38	123.08	4.8

CASCADES (CAS)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1872	12	14				48.3	120.3	7.4
1904	3	17	4	21	0	47.50	124.00	6.0
1945	4	29	20	16	17	47.40	121.70	5.5
1959	11	23	18	15	25	46.67	121.75	4.8
1961	9	16	3	24	58	46.00	122.20	4.3
1961	9	17	15	56	0	46.00	122.20	5.0
1961	10	31	2	35	0	48.40	120.00	4.3
1964	1	26	21	41	0	46.10	122.40	4.3
1966	8	17	14	39	50	48.20	125.00	3.8
1966	12	8	12	44	26	48.30	120.00	3.8
1969	10	9	17	7	58	46.90	121.60	4.4
1969	11	1	15	44	25	47.90	121.70	4.1
1969	11	10	07	38	45	48.55	121.51	4.3

CASCADES (CAS)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1971	11	23	2	12	19	48.32	121.44	4.1
1973	7	18	21	58	7	46.94	121.91	3.8
1974	4	20	3	0	9	46.76	121.52	4.8
1975	4	10	10	57	18	46.95	121.62	3.8
1975	11	30	10	48	21	49.23	123.62	4.9
1975	12	11	15	2	45	49.24	123.78	3.8
1976	1	18	8	38	11	48.52	124.63	3.9

NORTHERN VANCOUVER ISLAND (NVI)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1918	12	06	08	41	06	49.62	125.92	7.0
1927	05	07	22			50.15	127.85	5.5
1946	6	23	17	13	25	49.76	125.34	7.3
1957	12	16	17	27	48	49.82	126.48	6.0
1965	12	30	4	0	40	50.20	127.30	3.4
1966	1	13	7	49	6	49.67	126.82	4.0
1972	7	5	10	16	39	49.59	127.18	5.7
1974	2	12	3	4	53	50.21	127.61	3.4
1974	7	20	19	15	57	49.70	127.04	4.2
1974	9	20	11	33	49	50.10	127.79	3.5
1975	3	31	5	48	38	49.27	125.96	5.4
1975	11	29	10	50	30	49.43	126.79	4.4
1976	11	17	23	24	32	49.44	126.15	4.3
1978	05	25	21	53	44	50.20	127.70	3.3
1978	06	02	20	41	45	50.13	127.64	5.1
1978	06	03	11	54	40	50.19	127.60	4.6
1978	06	12	10	52	23	50.22	127.64	3.9
1978	06	13	11	39	30	50.23	127.53	3.6
1978	07	25	23	30	55	50.19	127.37	5.3

COAST MOUNTAINS (CSM)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1926	09	17	23	14	36	50.0	123.0	5.5
1942	01	31	06	49	11	51.18	123.58	5.5
1959	01	15	19	16	10	50.50	128.90	4.2
1959	12	29	12	7	15	52.30	127.80	3.8
1960	03	10	02	06	00	51.00	128.00	4.5
1961	7	24	10	39	24	50.60	128.90	3.9
1964	1	31	17	7	43	51.60	125.50	4.2
1964	5	10	13	44	2	51.40	129.20	4.1
1966	1	22	12	43	6	51.38	125.90	3.3
1966	6	10	5	47	50	51.00	125.20	3.4
1968	5	21	5	7	57	50.80	124.30	3.4
1968	6	18	5	37	57	51.10	129.00	4.1

COAST MOUNTAINS (CSM)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1968	11	1	10	24	59	51.00	124.20	4.5
1971	12	23	18	48	57	50.50	124.50	3.3
1976	8	12	6	28	60	50.64	123.05	3.8
1977	05	26	06	49	51	51.15	124.43	3.5

JUAN DE FUCA-EXPLORER (JFE)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1919	10	10	01	07	20	48.63	127.15	5.5
1920	3	29	5	7	53	50.50	129.50	6.4
1921	05	28	20	55		49.20	129.20	5.5
1924	3	30	0	8	56	50.50	129.50	6.0
1926	10	30	19	41	55	48.50	129.00	6.1
1926	11	1	1	39	18	48.75	128.50	6.6
1929	04	16	14	30	40	49.73	129.33	5.5
1929	09	17	19	17	39	50.92	129.58	6.3
1930	05	31	10	21	51	48.64	128.95	5.4
1932	08	18	20	23	00	48.39	127.62	5.5
1933	05	05	04	14	08	48.85	129.65	5.5
1935	09	24	22	12	20	49.63	129.17	6.2
1937	9	29	11	30	16	49.08	130.24	5.5
1938	04	22	04	15	54	50.03	128.45	5.5
1939	02	08	06	39	26	49.08	128.04	6.5
1939	07	18	03	26	38	49.01	129.22	6.5
1941	10	01	19	49	38	49.18	129.85	6.0
1941	11	06	17	31	54	49.35	129.83	6.0
1942	03	19	11	59	26	51.21	130.08	6.0
1942	6	9	11	6	45	49.20	129.90	5.7
1944	08	10	01	52	54	50.92	130.13	6.2
1944	08	13	08	22	28	50.13	130.46	5.8
1945	10	20	00	32	55	49.02	128.44	5.5
1946	7	17	22	7	0	50.00	129.00	6.5
1946	07	18	06	06	58	49.54	129.71	6.5
1946	11	12	14	35	44	49.10	128.40	5.5
1948	07	22	20	05	18	50.13	129.72	5.5
1948	07	22	20	52	43	49.84	129.65	5.5
1948	12	30	23	49	55	50.99	130.32	6.0
1951	09	27	19	24	13	49.45	128.60	5.8
1953	05	20	23	14	20	50.26	130.33	5.5
1953	12	04	14	54	48	49.41	129.02	6.3
1956	06	28	22	58	49	48.92	129.35	6.3
1957	03	24	08	22	23	50.85	130.36	6.0
1959	08	26	10	27	40	50.60	130.47	5.7
1960	4	14	0	37	52	48.50	130.40	5.7
1960	12	01	20	49	46	49.03	129.15	6.0
1961	02	01	00	36	01	50.26	129.63	5.5
1961	10	29	09	12	15	48.95	128.64	5.7
1962	06	02	12	26	07	50.00	129.65	5.7

JUAN DE FUCA-EXPLORER (JFE)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1964	03	31	09	01	29	50.83	130.05	5.7
1964	10	1	18	30	4	49.10	128.80	5.3
1965	2	26	15	43	17	50.30	129.80	4.5
1965	5	31	3	20	42	49.20	127.80	4.7
1965	7	2	22	55	56	49.20	129.80	3.9
1965	7	26	13	45	21	50.10	129.60	4.0
1965	8	12	9	4	37	50.20	129.70	4.9
1965	8	23	13	32	39	49.10	129.00	4.5
1965	8	31	1	9	15	50.50	129.00	3.9
1965	9	2	10	51	8	48.60	128.20	4.0
1965	9	2	11	37	50	48.60	128.00	4.6
1965	9	2	14	2	37	48.40	128.20	4.3
1965	9	2	15	42	26	48.30	128.40	4.4
1965	9	2	15	43	40	48.20	128.50	4.7
1965	9	2	18	1	20	48.30	128.30	4.4
1965	9	2	19	41	26	48.60	128.10	4.9
1965	9	2	21	16	44	48.60	128.00	4.0
1965	9	2	21	27	17	48.40	128.20	5.0
1965	9	3	0	30	31	48.50	128.00	4.0
1965	9	3	4	42	36	48.60	128.40	4.8
1965	9	11	7	13	19	50.30	129.50	4.9
1965	10	11	15	47	52	50.60	129.70	4.8
1965	10	11	17	54	48	50.60	129.90	4.2
1966	2	7	9	8	35	50.70	131.00	4.1
1966	2	7	13	45	42	50.60	131.50	3.8
1966	2	7	15	40	45	50.90	130.90	3.9
1966	2	8	7	58	20	50.90	130.30	3.9
1966	3	16	4	40	20	50.80	131.20	3.9
1966	3	30	12	39	56	49.80	129.90	5.1
1966	5	20	23	58	49	50.00	129.60	4.2
1966	9	1	14	11	21	49.30	129.30	4.6
1966	9	7	14	44	58	49.10	129.70	4.3
1966	9	9	18	33	52	49.20	129.40	4.8
1966	10	26	13	36	32	50.40	129.30	4.3
1966	11	4	20	30	9	48.90	128.90	4.2
1967	2	16	2	58	34	49.80	130.10	3.8
1967	3	5	11	11	2	51.20	129.50	3.8
1967	4	24	10	5	4	50.60	129.80	4.0
1967	4	29	0	4	42	51.10	130.10	5.3
1967	4	30	2	44	24	50.40	129.90	3.8
1967	8	27	13	34	51	50.20	130.00	4.3
1967	8	27	17	51	54	49.90	129.70	3.8
1967	8	27	18	29	5	50.10	129.80	3.9
1967	8	28	13	16	35	49.70	130.50	3.9
1967	8	28	13	49	40	49.80	129.70	3.8
1967	8	28	15	25	49	50.00	129.60	4.2
1967	8	28	16	20	4	50.20	129.70	4.1
1967	8	31	19	6	44	49.60	128.00	4.9
1967	10	16	13	27	33	49.20	129.30	4.9
1967	11	3	15	57	54	50.70	130.10	4.2

JUAN DE FUCA-EXPLORER (JFE)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1967	11	10	13	47	18	50.50	129.90	3.8
1967	12	9	18	31	40	49.20	128.70	4.0
1967	12	13	22	20	36	50.00	129.80	3.9
1968	2	1	3	5	1	50.20	130.40	4.2
1968	2	1	7	58	3	49.96	129.86	5.2
1968	2	15	18	27	30	51.35	129.68	3.8
1968	2	27	6	39	52	50.12	129.62	4.0
1968	3	2	3	14	45	49.25	128.92	4.5
1968	4	25	9	58	26	50.58	130.02	4.3
1968	6	18	5	37	54	50.87	130.10	3.8
1968	7	16	1	47	19	50.50	129.78	4.0
1968	7	28	21	16	49	50.53	129.70	4.0
1968	10	3	6	19	2	49.85	130.12	3.9
1968	11	17	21	11	34	49.00	128.90	4.4
1968	11	20	8	24	48	50.60	129.60	4.2
1968	11	22	11	59	26	49.00	128.70	4.0
1969	1	28	3	24	30	49.13	129.00	3.9
1969	3	10	22	50	47	50.50	129.60	4.1
1969	3	18	19	45	1	50.10	129.70	4.5
1969	3	18	20	31	28	50.17	129.88	5.1
1969	5	21	7	55	50	50.60	129.50	3.8
1969	5	21	9	20	27	50.68	129.50	3.9
1969	7	17	1	3	4	49.20	128.30	4.2
1969	9	4	13	22	58	49.40	129.20	4.2
1969	10	19	10	45	17	50.60	129.40	3.9
1969	10	23	21	36	41	50.40	129.90	4.1
1969	11	3	14	58	34	50.77	129.52	4.6
1970	2	1	23	2	31	50.00	129.00	4.0
1970	2	18	2	7	40	50.30	129.80	4.7
1970	7	23	13	31	40	48.50	128.00	3.8
1970	11	10	2	10	43	50.60	129.50	4.1
1970	11	16	12	49	21	49.30	128.10	4.5
1970	12	31	1	27	8	50.20	129.50	4.3
1970	12	31	5	34	14	47.78	128.77	5.2
1970	12	31	10	46	16	47.80	128.20	4.2
1971	1	1	6	50	53	47.80	128.60	4.5
1971	3	10	15	38	26	49.35	127.46	5.0
1971	3	13	23	51	38	50.56	129.90	6.4
1971	3	14	0	10	39	50.60	129.90	4.3
1971	3	14	0	44	16	50.60	129.90	4.0
1971	3	14	0	51	7	50.60	129.90	4.3
1971	3	14	4	41	58	50.60	129.90	4.1
1971	3	15	5	18	56	50.60	129.90	4.2
1971	6	29	6	28	54	51.20	129.60	4.2
1971	7	22	14	5	30	50.60	129.20	4.2
1971	11	20	21	24	37	48.60	129.40	5.0
1971	11	25	23	1	33	49.10	129.00	4.4
1971	11	25	23	40	8	49.00	129.10	5.1
1971	12	5	5	50	8	49.40	129.50	5.2
1971	12	5	6	12	51	49.80	129.20	5.0

JUAN DE FUCA-EXPLORER (JFE)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1971	12	8	8	25	2	49.20	128.10	4.6
1971	12	8	8	35	25	49.20	128.10	5.0
1971	12	10	20	25	14	49.80	129.00	4.0
1971	12	11	10	39	11	49.20	128.40	4.2
1971	12	20	9	15	8	49.20	128.50	3.8
1971	12	25	18	18	35	49.30	128.50	4.3
1971	12	30	7	45	11	49.10	128.80	4.0
1972	1	14	22	23	43	50.32	130.79	4.4
1972	5	14	20	35	41	50.55	130.47	3.9
1972	7	23	10	52	33	50.11	129.09	4.8
1972	7	23	19	13	9	50.10	129.30	5.8
1972	7	23	21	43	5	50.06	129.30	4.8
1972	7	23	20	17	32	50.21	129.16	4.8
1972	7	23	21	43	5	50.06	129.30	4.8
1972	11	1	16	11	35	49.93	130.12	3.9
1973	3	28	6	23	7	51.19	129.69	3.9
1973	4	17	2	16	6	50.71	130.65	3.8
1973	6	3	7	23	29	50.55	130.31	4.2
1973	7	13	2	59	30	49.12	127.84	4.8
1973	7	13	2	59	39	49.02	128.02	4.5
1974	1	29	6	13	11	49.43	129.04	3.8
1974	3	7	7	50	26	50.59	130.36	3.9
1974	5	30	1	0	2	49.28	127.65	3.8
1974	8	1	22	10	43	50.64	130.35	3.9
1975	1	29	16	16	56	50.06	129.99	3.8
1975	1	29	17	43	10	49.97	130.14	3.8
1975	2	18	20	21	9	50.75	130.55	3.8
1975	2	18	21	6	48	50.71	129.77	4.0
1975	2	21	16	43	28	50.81	130.57	3.8
1975	3	20	20	36	54	50.54	130.25	4.1
1975	8	1	14	4	26	49.27	128.96	3.8
1975	11	24	10	35	46	50.51	130.49	4.0
1975	12	11	6	28	34	50.05	130.19	4.1
1975	12	11	7	3	14	50.09	130.11	3.9
1975	12	12	1	48	41	49.68	130.16	4.1
1975	12	12	1	52	40	50.02	130.23	4.3
1975	12	12	2	14	32	49.98	130.24	4.1
1976	1	1	4	11	43	50.19	130.18	4.3
1976	1	2	3	36	21	50.38	130.02	4.4
1976	2	27	13	8	18	50.87	130.71	4.0
1976	2	28	0	40	5	50.96	130.64	3.8
1976	4	25	11	20	11	49.41	127.11	4.4
1976	6	6	2	17	18	49.04	127.86	5.0
1976	8	26	6	43	10	50.61	130.26	3.8
1976	11	9	20	17	19	50.61	129.84	4.4
1976	12	20	17	12	45	49.00	128.88	4.7
1976	12	20	20	33	8	48.80	129.29	6.7
1976	12	20	21	6	39	48.90	128.72	5.1
1976	12	20	21	12	52	49.19	129.17	4.1
1976	12	20	21	21	37	48.94	128.41	3.8

JUAN DE FUCA-EXPLORER (JFE)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1976	12	26	10	52	48	49.38	130.03	3.8
1977	04	03	18	45	46	50.56	130.18	4.0
1977	07	09	08	04	18	50.86	130.72	4.2
1977	12	25	07	21	01	50.61	129.98	4.1

QUEEN CHARLOTTE FAULT (QCF)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1917	7	1	13	20	50	50.00	128.00	6.4
1919	5	18	10	23	56	56.00	136.00	6.0
1921	04	10	13	40	16	54.42	133.66	6.5
1929	05	26	22	40	01	51.19	130.16	7.0
1938	03	22	15	22	20	52.54	131.90	6.3
1940	9	29	5	57	12	54.00	134.00	5.5
1945	10	29	10	54	17	51.59	130.98	5.5
1948	02	28	01	58	05	53.41	132.74	6.5
1949	08	22	04	01	12	53.62	133.27	8.0
1949	8	23	19	43	34	52.49	132.65	5.5
1949	08	23	20	24	31	52.69	132.23	6.4
1949	08	24	22	37	13	52.78	132.11	5.5
1949	8	26	22	39	29	54.50	136.00	5.5
1949	10	31	1	39	28	56.00	136.00	6.7
1950	05	22	19	49	43	51.56	130.51	5.7
1950	9	28	21	47	1	54.50	134.50	5.5
1956	02	19	02	18	00	51.61	131.37	6.8
1956	11	17	20	27	15	54.50	134.00	6.5
1956	12	21	08	58	55	51.29	130.60	6.7
1959	01	16	16	50	43	51.98	131.21	5.4
1960	07	04	04	28	35	51.79	131.19	6.6
1960	07	04	13	10	07	51.79	131.09	6.0
1965	3	5	9	39	24	51.50	131.10	4.0
1965	3	5	12	40	39	52.70	132.00	3.9
1965	6	28	23	15	8	51.00	133.00	4.1
1966	1	20	19	51	14	51.30	131.20	4.2
1966	1	20	19	59	42	51.30	132.10	3.8
1966	1	21	11	27	52	51.30	131.50	3.8
1966	2	7	14	12	26	50.90	131.20	4.0
1966	2	7	14	32	21	51.30	130.70	3.8
1966	3	3	7	11	55	51.60	131.50	3.8
1966	3	25	21	59	26	56.60	135.40	4.7
1966	4	16	22	49	39	56.70	136.20	4.1
1966	4	17	16	46	47	54.10	133.60	5.0
1966	7	16	6	22	43	52.20	133.60	3.8
1966	7	23	19	34	58	54.20	135.20	4.3
1966	12	8	9	33	48	51.30	130.60	3.9
1967	4	12	0	54	42	56.20	136.00	4.4
1967	4	19	18	12	31	52.50	132.10	4.6
1968	1	15	12	24	14	52.20	131.60	4.3

QUEEN CHARLOTTE FAULT (QCF)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1968	1	15	14	29	20	52.20	131.60	3.9
1968	6	13	8	51	16	51.40	131.55	4.5
1968	9	22	3	51	51	51.38	131.22	4.4
1968	11	16	14	3	22	56.20	138.70	3.8
1969	12	15	8	31	30	53.00	135.00	4.0
1970	2	18	9	23	2	52.40	131.50	4.8
1970	6	24	7	30	28	51.86	130.97	4.9
1970	06	24	13	09	08	51.77	130.76	7.0
1970	6	24	13	17	4	51.74	131.20	5.2
1970	7	3	6	20	10	51.80	131.00	4.0
1971	2	5	7	33	28	51.75	130.65	5.5
1971	2	11	6	24	46	54.70	135.20	4.6
1971	5	28	12	11	3	52.40	132.40	4.6
1971	11	6	11	12	52	51.15	135.28	4.0
1971	12	9	20	33	15	56.43	135.59	4.2
1972	6	17	23	50	23	54.27	133.61	4.3
1972	6	18	20	43	12	54.32	133.62	3.8
1972	7	30	21	45	16	56.77	135.91	7.6
1972	7	30	22	51	54	56.30	136.10	4.5
1972	7	31	3	13	55	56.63	135.78	4.3
1972	7	31	9	47	14	56.01	135.53	4.2
1972	7	31	11	25	33	56.67	135.76	3.9
1972	8	4	9	48	11	56.23	135.64	4.9
1972	8	4	11	38	8	56.19	135.42	5.5
1972	8	7	8	31	45	55.99	136.40	3.8
1972	8	9	19	20	46	56.23	135.61	3.9
1972	8	10	21	39	52	55.98	135.67	4.0
1972	8	15	10	56	13	56.31	135.57	5.4
1972	11	7	17	33	29	56.23	135.71	4.4
1972	11	15	9	11	38	56.34	136.15	4.2
1972	11	17	16	41	35	56.03	135.60	4.8
1972	12	8	18	56	56	56.30	135.90	4.2
1973	1	12	5	59	32	56.10	135.51	4.3
1973	1	14	1	23	7	56.27	135.52	4.3
1973	9	5	1	14	3	53.87	133.07	4.0
1973	9	6	17	15	48	51.17	130.40	4.3
1974	4	8	23	24	41	54.13	134.10	4.0
1974	8	20	4	25	4	51.06	130.70	4.3
1975	2	14	12	15	6	52.68	132.04	3.8
1975	4	7	1	47	45	51.64	130.88	4.7
1975	5	23	15	12	30	51.22	131.23	4.7
1975	7	14	18	19	25	51.46	131.83	3.9
1975	8	20	2	6	1	51.22	130.76	4.5
1976	2	23	12	12	31	51.46	130.76	4.1
1976	2	23	14	58	59	51.60	130.51	4.3
1976	2	23	15	14	16	51.50	130.50	6.0
1976	2	23	16	12	23	51.47	130.68	4.7
1976	2	23	18	12	35	51.43	130.73	4.5
1976	2	27	10	39	31	51.07	130.57	3.8
1976	2	28	1	7	17	51.00	130.76	4.1

QUEEN CHARLOTTE FAULT (QCF)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1976	5	13	7	11	44	52.86	132.24	4.8
1976	6	18	9	19	24	53.92	133.06	4.3
1976	7	17	23	40	1	53.91	132.94	3.9
1976	10	15	20	29	33	54.31	133.67	3.8
1976	11	10	19	16	34	51.37	130.67	4.1
1977	01	14	14	55	58	51.34	130.33	3.8
1977	08	08	14	23	58	55.56	135.26	4.6

SANDSPIT FAULT (SPT)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1929	03	01	07	31	14	51.79	129.74	6.1
1936	12	21	19	03	16	53.02	131.65	6.0
1945	08	02	20	44	45	53.89	132.63	6.2
1965	5	15	3	14	32	55.00	133.50	4.1
1965	6	5	8	38	56	54.00	133.00	4.1
1966	6	22	16	55	24	52.40	130.10	3.9
1967	2	22	17	9	25	52.40	131.20	3.8
1968	6	28	18	4	42	56.00	133.60	3.9
1970	2	19	8	9	18	53.30	132.30	4.0
1970	6	24	17	16	53	51.90	130.40	3.9
1970	6	24	19	10	19	51.90	130.20	3.9
1971	7	15	0	24	2	54.60	133.60	5.3
1974	4	3	22	46	33	54.12	133.41	4.5
1974	4	8	23	24	41	54.25	133.36	4.5
1976	2	23	12	12	35	51.60	130.10	3.8
1976	7	17	23	40	1	53.91	132.94	3.9
1977	07	19	17	22	10	54.17	133.17	4.6

SOUTHEASTERN B.C. (SBC)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1918	02	04	20	37		52.28	118.37	6.0
1942	11	01	18	50	06	48.00	116.70	5.5
1962	08	28	19	19	59	51.7	121.9	4.3
1965	03	23	00	28	18	49.7	117.4	4.0
1965	03	24	12	44	55	49.9	117.4	3.8
1965	04	21	11	47	30	52.3	117.2	3.7
1965	04	28	19	00		48.6	116.9	4.3
1965	07	28	18	33	45	52.3	119.8	3.5
1966	01	02	10	10	51	51.5	116.5	4.5
1966	05	29	13	47	27	49.4	114.9	3.5
1966	11	06	10	50	54	48.3	119.5	3.8
1967	04	05	02	55	25	48.2	119.1	3.4
1967	04	20	15	19	33	48.2	119.42	3.3
1967	06	04	03	26	23	52.2	120.0	3.4

SOUTHEASTERN B.C. (SBC)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1967	06	06	17	12	56	48.2	119.4	3.3
1968	04	12	10	26	08	48.6	116.17	3.6
1968	07	14	03	32	36	50.58	117.50	3.7
1968	07	26	22	23	30	52.28	118.68	3.3
1968	08	31	08	31	18	49.42	116.92	3.7
1968	12	13	08	50	08	52.40	120.20	3.7
1969	05	10	17	48	49	49.14	118.77	3.4
1970	05	30	19	36	51	49.20	113.70	3.7
1970	11	27	22	17	50	52.64	119.13	3.8
1973	03	04	02	47	32	52.06	118.07	3.9
1973	03	04	05	02	43	52.03	118.04	3.5
1973	03	16	06	28	56	48.91	114.80	3.5
1973	03	22	21	21	51	52.06	118.01	3.9
1974	07	26	23	36	03	48.72	114.89	3.7
1977	06	12	02	57	06	51.54	118.46	3.6
1977	10	09	16	42	39	53.65	118.29	4.4

NORTHERN B.C. (NBC)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1965	09	05	10	04	17	55.0	130.6	4.5
1969	10	20	01	48	55	57.3	126.6	4.4
1971	10	12	19	09	53	59.78	135.29	3.4
1973	11	05	12	36	17	54.43	129.06	4.2
1973	11	06	15	57	12	54.46	128.93	3.7
1974	02	18	03	53	25	57.22	124.92	3.5
1975	06	05	05	52	40	57.95	124.80	4.2
1975	07	11	01	14	10	58.47	133.38	3.3
1976	04	07	01	53	34	57.64	125.27	3.5
1976	07	11	12	58	09	58.45	133.38	3.3
1977	09	29	22	19	23	54.83	126.97	3.3

SOUTHERN SASKATCHEWAN (SAS)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1909	05	16	04	15		49.	104.	5.5
1943	06	25	04	25		48.5	105.0	4.0
1968	10	11	12	28	04	49.61	104.49	2.8
1969	10	06	20	24	53	48.29	106.58	3.1
1972	07	26	03	58	19	49.35	104.93	3.7
1973	09	26	18	38	27	47.12	106.13	2.8
1975	09	05	20	47	41	48.36	104.38	3.5
1976	03	23	22	31	47	49.56	104.37	3.2
1976	03	25	00	12	16	49.39	104.27	3.5

FAIRWEATHER-YAKUTAT (FWY)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1899	9	4	0	22	0	60.00	142.00	8.2
1899	9	10	17	4	0	60.00	140.00	7.8
1899	9	10	21	41	0	60.00	140.00	8.6
1900	10	9	12	28	0	60.00	142.00	8.2
1927	10	24	15	59	55	57.50	137.00	7.1
1947	4	30	4	49	46	59.00	139.00	6.3
1957	6	23	3	27	2	58.50	137.00	5.6
1958	7	10	6	15	51	58.60	137.10	7.9
1958	7	13	8	10	2	58.30	136.90	5.6
1958	9	24	3	44	14	59.50	143.50	6.2
1963	6	17	18	32	14	60.50	140.80	5.4
1965	04	26	01	57	14	58.90	142.70	5.3
1965	04	28	09	40	19	58.60	143.30	4.9
1965	06	07	11	24	49	60.30	140.90	4.8
1965	6	27	11	8	56	60.28	141.13	5.1
1965	09	30	23	47	40	59.70	143.40	5.0
1965	10	12	08	16	23	59.50	144.60	4.8
1965	12	23	20	47	36	60.60	140.70	5.8
1966	01	15	11	59	58	59.50	144.60	5.3
1966	08	07	14	11	55	59.60	144.60	4.8
1966	10	10	21	17	35	57.40	136.10	4.8
1967	05	17	00	33	12	60.80	143.65	4.8
1969	06	02	09	47	59	59.45	144.67	4.7
1969	06	11	00	58	10	59.60	144.80	5.3
1969	06	11	01	05	03	59.59	144.76	4.9
1970	01	10	04	21	43	59.45	144.97	4.9
1970	02	24	08	05	40	59.57	143.87	5.0
1970	04	11	04	05	41	59.70	142.70	6.2
1970	04	11	09	59	46	59.51	142.75	4.6
1970	04	11	12	55	38	59.77	142.66	4.5
1970	04	16	05	33	17	59.80	142.60	6.2
1970	04	17	15	31	47	59.55	142.70	4.3
1970	04	19	01	15	47	59.60	142.80	6.0
1970	04	21	06	44	25	59.62	142.65	4.9
1970	8	21	11	58	50	60.77	142.60	4.5
1970	9	6	15	43	19	60.17	141.14	4.7
1971	1	1	4	45	30	59.62	144.65	5.1
1971	1	2	19	9	51	59.58	144.70	4.8
1971	3	26	17	35	17	60.33	140.94	5.8
1971	3	27	22	0	5	60.05	140.92	4.5
1972	3	12	5	56	17	58.98	137.90	4.3
1972	4	17	2	29	50	60.81	144.61	3.8
1972	5	29	12	8	12	57.40	137.50	4.2
1972	8	13	2	30	36	59.31	144.90	4.6
1972	8	18	22	4	45	57.40	136.10	4.1
1972	9	29	9	0	36	60.21	141.03	4.7
1972	12	7	22	19	44	57.09	136.52	3.9
1973	2	7	15	26	44	59.40	143.32	4.5
1973	6	26	0	41	49	59.41	144.69	4.4
1973	7	1	13	33	35	57.84	137.33	6.1

FAIRWEATHER-YAKUTAT (FWY)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1973	7	1	15	6	37	57.62	137.74	3.8
1973	7	1	15	12	5	57.78	137.28	5.2
1973	7	1	16	3	10	57.68	137.50	4.2
1973	7	2	13	36	52	57.93	136.92	3.8
1973	7	2	22	54	45	57.93	137.69	4.5
1973	7	3	16	30	37	58.05	137.73	4.6
1973	7	3	16	59	35	57.98	138.02	6.0
1973	7	3	17	44	16	57.99	137.88	5.1
1973	7	4	7	17	6	58.06	137.31	4.5
1973	7	4	13	26	20	58.01	137.85	4.6
1973	7	5	7	45	24	58.01	137.29	4.5
1973	7	5	7	49	4	57.90	137.90	5.4
1973	7	5	8	51	30	58.03	137.37	3.9
1973	7	11	23	16	27	57.92	138.06	4.6
1973	7	13	5	11	5	60.08	140.89	3.9
1973	7	14	5	8	22	58.00	138.00	5.0
1973	7	16	21	20	16	57.69	137.60	3.8
1973	7	27	13	54	50	58.05	137.66	4.0
1973	7	28	19	58	47	58.00	137.89	4.7
1973	8	9	6	28	24	57.83	137.39	3.8
1973	9	9	2	54	47	60.25	140.80	3.8
1973	9	12	7	0	23	60.15	140.85	3.9
1974	2	21	16	28	4	60.33	140.59	4.1
1974	2	23	8	9	46	60.66	142.86	3.9
1974	3	4	6	54	33	60.11	140.67	3.9
1974	3	4	6	54	33	60.11	140.67	3.9
1974	4	18	21	54	26	59.16	139.97	3.9
1974	7	21	16	33	34	59.13	140.27	4.1
1974	08	28	18	43	26	59.51	144.45	4.9
1974	9	20	1	49	37	59.96	141.45	3.9
1974	9	28	17	33	33	60.05	140.62	4.1
1974	11	5	10	24	54	60.05	140.43	4.0
1975	4	30	20	9	55	57.26	135.79	3.8
1975	9	24	14	17	54	59.88	141.85	4.2
1976	1	25	18	52	41	59.89	141.55	4.5
1976	2	15	21	15	49	57.91	138.57	4.1
1976	4	20	14	27	21	60.28	140.58	4.5
1976	7	2	16	28	40	60.52	141.24	4.0
1976	7	8	3	59	49	60.32	140.89	3.8
1976	11	7	14	48	48	57.88	137.79	4.2
1976	11	12	18	48	37	59.61	138.95	3.8
1977	01	13	22	05	59	59.43	142.23	4.5
1977	01	20	15	45	26	57.86	137.86	4.3
1977	10	30	10	51	58	59.79	141.32	3.8

DENALI-SHAKWAK (DSK)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1944	02	03	12	14	59	60.10	137.88	6.5
1952	03	09	20	00	17	59.10	136.70	6.0
1958	08	31	23	00	16	63.30	144.20	5.9
1968	2	16	2	42	34	61.19	139.99	4.5
1971	4	24	0	39	18	60.53	139.05	4.5
1972	06	10	03	31	24	61.52	140.21	5.3
1972	6	10	9	46	19	61.37	140.12	3.8
1972	6	11	1	11	57	61.51	139.91	3.8
1973	9	30	17	33	50	61.71	141.21	4.1
1974	1	27	4	39	38	59.35	136.37	4.0
1974	2	7	13	51	55	59.17	137.03	4.0
1974	04	15	16	27	36	59.19	136.43	4.2
1974	08	01	02	02	30	60.86	137.99	4.5
1975	2	17	0	38	3	60.78	139.76	3.8
1977	01	28	23	29	02	59.01	136.82	3.8
1977	11	06	19	11	03	62.09	144.87	3.8

RICHARDSON MTS (RIC)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1940	6	5	11	1	10	66.90	135.00	6.5
1940	6	5	11	1	10	66.90	135.00	6.5
1952	6	15	15	12	40	65.60	134.90	5.5
1953	1	11	22	53	30	65.30	133.20	5.5
1955	3	1	4	42	59	65.30	132.80	6.6
1956	1	7	16	41	4	65.60	133.70	5.5
1957	12	9	22	7	43	65.30	133.50	5.7
1965	10	5	0	17	14	65.40	133.00	5.2
1968	1	27	18	17	54	66.03	135.02	4.0
1968	4	26	15	49	26	65.37	133.28	4.2
1971	3	4	13	43	11	66.74	135.37	4.1
1971	11	21	13	24	57	65.92	134.94	4.2
1972	2	11	8	15	12	65.20	133.30	4.0
1972	2	18	6	33	6	64.89	133.21	4.0
1972	7	1	17	6	53	65.44	133.55	3.8
1972	7	26	18	46	22	66.52	135.97	4.6
1973	2	16	8	34	22	66.87	135.49	4.3
1974	5	12	18	15	13	65.41	133.84	3.9
1974	12	9	4	27	7	64.97	133.51	4.0
1975	6	25	5	44	7	64.75	133.56	3.8
1976	1	16	12	37	18	65.39	134.25	4.2
1976	2	19	4	55	42	66.41	135.28	5.0
1976	4	8	18	59	40	64.93	134.03	4.2
1976	11	3	10	29	20	65.16	133.55	4.2
1976	11	28	1	23	41	66.38	135.24	4.2
1977	10	13	22	02	04	65.56	135.06	4.0

BEAUFORT (BFT)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1968	4	4	0	33	30	71.92	132.68	3.8
1968	8	6	9	9	54	72.39	136.33	4.4
1970	5	4	15	14	59	72.10	132.95	4.0
1971	9	27	21	49	27	70.95	131.96	4.3
1972	12	15	9	8	35	71.64	134.96	4.5
1975	4	6	19	25	36	71.55	133.02	4.3
1975	6	14	20	50	26	71.96	131.72	5.1
1975	12	8	13	36	3	71.02	132.90	4.2
1976	6	30	6	29	54	72.13	135.55	4.2
1976	11	12	6	40	41	71.54	135.86	4.9

MACKENZIE (MKZ)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1968	1	22	9	51	24	70.20	144.28	3.8
1968	1	22	14	4	52	70.35	143.88	4.3
1968	1	22	23	44	34	70.41	143.77	5.3
1968	1	23	8	30	45	70.40	144.24	4.1
1968	1	23	20	57	52	70.36	144.29	4.3
1968	1	30	9	28	33	70.25	144.32	3.8
1968	2	5	4	7	21	70.32	144.24	3.9
1968	2	6	16	36	22	70.36	143.91	4.5
1968	2	6	18	42	54	70.39	144.13	3.9
1968	2	10	17	29	0	70.34	143.89	4.3
1968	2	10	17	29	19	70.23	143.63	4.3
1968	2	10	17	39	50	70.54	142.65	3.9
1968	2	13	0	59	2	70.42	143.27	4.1
1968	2	28	8	36	16	70.41	143.16	4.1
1968	3	9	13	55	37	70.27	144.10	4.2
1968	4	25	10	33	50	70.21	144.46	4.4
1969	3	13	18	43	50	63.56	128.36	5.0
1969	7	3	21	30	26	66.03	140.26	3.8
1969	10	20	1	48	55	57.30	126.60	4.4
1969	10	28	0	46	4	68.03	136.46	4.0
1970	5	19	11	32	53	69.24	130.59	4.0
1970	12	19	15	24	16	64.60	138.94	3.8
1971	1	28	5	6	44	66.59	141.44	4.2
1971	3	2	0	32	5	70.23	144.07	4.3
1971	3	29	16	39	8	62.82	123.39	4.0
1971	12	14	15	18	31	66.58	139.46	3.8
1972	2	12	17	52	35	62.62	124.84	4.9
1973	3	23	15	15	37	66.69	130.21	4.6
1974	8	14	19	55	39	62.43	124.45	4.1
1974	9	22	11	7	50	65.22	141.26	4.3
1975	3	5	9	46	52	63.35	130.09	4.1
1975	3	30	11	33	34	69.60	143.23	3.9
1975	3	31	12	53	3	69.98	142.54	3.8
1975	6	5	5	52	40	57.95	124.80	4.2

MACKENZIE (MKZ)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1975	8	24	0	27	47	62.54	127.76	4.3
1975	9	3	1	23	54	64.66	138.51	4.7
1976	2	25	18	15	28	65.46	139.99	3.8
1976	3	5	3	18	20	71.21	142.09	4.2
1976	3	17	23	59	58	67.05	131.76	4.3
1976	8	8	19	55	27	64.40	137.46	4.2
1976	8	16	13	36	54	64.77	134.87	4.1
1977	01	06	12	51	28	69.54	129.70	4.2
1977	07	03	17	41	16	62.54	128.69	4.0

CHARLEVOIX (CHV)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1663	2	5	17	30		47.60	70.10	7.0
1860	10	17	11	15		47.50	70.10	6.0
1870	10	20	16	30		47.40	70.50	6.5
1924	9	30	8	52	30	47.80	69.80	5.5
1925	3	1	2	19	20	47.80	69.80	7.0
1925	3	1	4	30	42	47.80	69.80	5.0
1925	3	21	15	22	4	47.80	69.80	5.0
1930	12	25	22	7	34	47.30	70.40	4.5
1931	1	8		13	36	47.30	70.40	5.5
1939	6	24	19	20	21	47.30	70.40	4.5
1939	10	19	11	53	58	47.80	69.80	5.0
1939	10	21	8	7	13	47.80	69.80	4.0
1939	10	27	1	36	36	47.80	69.80	4.5
1939	11	7	2	40	32	47.80	70.50	4.0
1939	12	25	10	29	13	48.10	70.40	4.0
1940	4	13	8	13	34	47.73	70.73	3.8
1940	10	13	19	50	51	47.80	69.80	4.5
1941	9	6	17	4	56	47.43	70.52	3.8
1941	10	6	16	34	27	47.63	70.60	4.0
1943	9	28	16	30	25	47.27	70.40	3.8
1943	11	6		6	40	47.60	70.30	3.9
1944	2	5	12	37	52	47.40	70.50	4.0
1945	10	9	13	18	44	47.80	69.80	4.5
1947	2	2	16	50	32	47.67	70.53	4.0
1947	3	29	12	28	52	47.37	70.50	4.0
1947	10	22	9	36	38	47.55	70.72	3.8
1948	1	1	18	33	45	47.30	70.40	4.5
1952	3	30	13	11	7	47.60	69.88	4.0
1952	4	19	2	50	52	47.47	70.58	3.8
1952	10	14	22	3	42	47.80	69.80	5.0
1954	2	7	20	24	16	47.60	70.25	3.8
1955	2	1	12	40	27	47.50	70.30	4.0
1957	8	6	23	50	38	47.30	70.42	4.0
1958	8	8	22	15	3	47.93	70.38	3.9
1960	4	23	11	47	52	47.53	70.30	4.0

CHARLEVOIX (CHV)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1962	7	27	17	56	57	47.25	70.67	4.0
1965	12	16	13	53	19	47.50	69.90	4.0
1968	3	30	15	28	59	47.94	70.49	3.1
1968	4	11	9	18	33	47.60	70.44	3.5
1968	10	20	2	36	58	47.47	70.57	3.6
1969	5	10	18	43	29	47.47	70.65	3.6
1969	5	10	20	1	53	47.47	70.65	3.6
1969	7	14	3	6	59	47.83	70.09	4.0
1969	8	31	7	20	27	47.49	70.07	3.2
1970	9	7	21	39	27	47.92	70.30	3.2
1971	9	12	8	31	43	47.56	70.24	3.2
1972	2	13	11	8	7	47.77	70.27	3.2
1972	8	2	1	3	1	47.40	70.50	2.9
1972	9	25	11	30	21	47.50	70.60	3.0
1973	1	28	13	7	50	47.98	70.00	3.1
1973	9	10	6	11	12	47.68	70.24	2.9
1973	11	16	1	36	34	47.55	70.29	3.1
1974	6	30	16	55	10	47.84	70.08	3.1
1975	08	21	04	29	37	47.44	70.18	3.1
1975	11	25	23	29	14	47.62	70.09	2.9
1976	05	20	14	55	16	47.45	70.31	2.8
1976	07	11	05	15	02	47.43	70.44	2.9
1976	08	03	02	57	13	47.69	70.10	2.9
1976	10	23	20	58	18	47.82	69.78	4.2
1976	10	23	21	23	06	47.88	69.78	3.1
1976	10	24	10	49	46	47.81	69.87	3.5
1977	02	14	00	35	04	47.54	70.42	3.1
1977	06	20	05	05	53	47.84	70.16	3.1

WESTERN QUEBEC (WQU)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1914	2	10	18	31		46.00	75.00	5.5
1931	9	23	22	47	37	47.00	76.07	4.5
1934	4	15	2	58	13	44.90	73.90	4.5
1935	11	1	6	3	40	46.78	79.07	6.0
1935	11	2	14	31	58	47.70	78.30	5.0
1937	11	6	14	31	20	46.73	75.72	4.0
1937	11	12	16	57	32	45.92	74.33	4.0
1938	11	18	22	19	6	44.75	75.25	4.0
1938	11	26	7	47	57	47.03	76.20	4.0
1938	12	25	7	46	19	47.58	75.37	4.0
1940	2	10	20	57	17	46.30	76.30	4.0
1941	6	26	4	5	44	47.40	76.83	4.0
1942	5	20	12	19	22	45.90	74.67	4.5
1942	5	24	11	33	57	44.73	73.83	3.9
1942	12	5	21	10	51	46.97	76.07	4.0
1943	7	6	22	10	14	44.92	73.13	4.0

WESTERN QUEBEC (WQU)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1944	1	22	21	55	9	45.70	76.60	4.5
1944	3	8	12	49	56	46.68	78.87	4.0
1944	9	5	4	38	45	44.97	74.90	5.6
1944	9	5	8	51	6	44.98	74.90	4.5
1944	9	9	23	24	48	44.98	74.90	4.0
1944	10	31	8	42	25	44.98	74.90	4.0
1945	6	12	7	58	15	46.90	75.50	4.5
1947	1	19		45	1	46.80	76.70	4.0
1948	5	7	12	2	26	45.75	73.63	4.0
1949	10	16	23	33	42	45.30	74.83	4.0
1950	3	6	16	14	11	46.00	74.50	4.0
1950	4	14	18	20	48	48.00	75.70	5.0
1950	8	4	14	29	28	45.20	74.72	4.0
1951	10	25	7	7	52	45.10	74.73	4.0
1952	1	30	4			44.50	73.20	4.5
1952	3	17	4	14	41	47.30	76.40	4.0
1952	7	19	1	16	17	46.87	75.83	4.5
1954	4	12	21	22	1	46.90	76.05	4.5
1954	9	11	18	55	52	47.33	75.63	4.5
1955	2	3	2	30		44.50	73.22	4.0
1956	6	15		53	37	47.10	76.43	4.0
1956	11	4	11	53	24	46.22	75.73	4.0
1958	3	1	17	41	49	46.90	76.03	3.9
1958	5	14	17	41	21	46.97	76.55	5.0
1958	7	25	3	45	11	46.57	75.80	3.8
1959	5	21	9	38	51	46.55	76.45	4.0
1962	1	27	12	11	17	45.92	74.85	4.0
1963	8	26	16	29	35	45.18	73.95	3.5
1963	10	15	12	29	2	46.35	77.59	4.0
1963	10	15	13	59	53	46.30	77.59	4.0
1964	1	8	8	59	28	46.23	77.53	3.3
1964	1	8	10	3	26	46.23	77.53	4.0
1964	1	8	10	4	31	46.23	77.53	4.5
1964	3	29	4	16		44.90	74.90	4.0
1964	7	24	10	34	11	46.65	76.25	3.3
1965	9	15	17	56	28	46.72	79.05	3.8
1965	11	7	20	57	44	47.30	76.20	4.0
1965	11	24	21	28	1	46.93	76.28	3.7
1965	12	19	1	5	52	47.03	76.42	3.5
1966	6	25		5	51	45.16	73.83	3.4
1966	11	13	15	43	29	47.00	76.25	3.6
1967	6	11	1	49	39	46.58	75.03	3.7
1968	10	19	10	37	18	45.30	74.12	3.2
1968	11	3	20	50	49	46.17	76.30	3.1
1969	3	19	7		37	45.64	76.22	2.8
1969	6	12	11		11	46.92	75.95	2.9
1969	10	10		7	7	46.42	75.20	4.0
1969	10	10	8	16	12	46.38	75.05	2.8
1970	4	6	11	29	16	46.16	74.84	2.8
1970	10	15	18	56	11	47.07	76.25	3.3

WESTERN QUEBEC (WQU)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1971	1	6	6	22	8	47.17	75.96	3.0
1971	1	19	13	44	25	46.92	75.18	3.1
1971	5	14	6	20	9	45.10	73.37	3.2
1971	7	6	17	47	49	46.55	76.28	3.0
1971	9	27	8	47	23	45.71	75.17	3.2
1971	11	15	10	38	55	45.06	73.87	3.0
1971	11	22	5	29	7	47.24	76.28	3.0
1971	11	23	16	32	30	45.83	76.62	3.0
1971	12	18	15	36	24	46.18	74.62	4.0
1972	4	25	3	24	25	46.60	76.00	3.5
1972	6	2	4	24	57	45.70	75.90	2.8
1972	7	30	10	42	16	46.30	76.10	3.1
1972	9	12	9	15	40	46.10	77.60	3.1
1972	12	16	19	1	36	45.77	75.22	4.0
1973	2	2	23	9	30	44.43	74.78	2.8
1973	2	25	19	46	46	45.23	73.97	2.9
1974	2	13	18	14	53	46.40	75.27	2.9
1974	3	18	16	5		44.45	74.85	3.0
1974	4	29	6	10	48	46.00	75.23	2.8
1974	8	8	11	55	33	45.93	76.08	3.2
1974	10	23	22	52	57	46.08	75.48	3.2
1974	11	2	13	47	56	46.07	75.03	3.2
1974	11	3	4	27	4	46.07	75.05	2.8
1974	12	2	10	58	5	46.25	75.50	3.5
1975	04	03	19	03	17	45.73	74.24	3.1
1975	05	29	21	19	16	47.23	75.19	3.2
1975	06	09	18	39	22	44.94	73.65	3.5
1975	07	12	12	37	15	46.46	76.22	4.2
1975	12	19	15	25	11	47.01	78.84	3.8
1976	01	13	21	15	58	46.88	76.09	2.9
1976	02	02	14	44	13	46.10	75.56	2.9
1976	07	13	03	51	14	45.17	74.10	3.1
1976	11	05	16	50	00	46.76	75.48	2.9
1976	11	06	06	09	29	47.11	75.96	3.0
1977	01	08	05	05	23	47.25	75.55	2.9
1977	07	14	07	39	30	45.98	74.41	3.4
1977	09	28	17	21	44	44.45	73.92	2.9
1977	11	07	20	48	52	46.29	75.21	3.0
1977	11	25	18	47	22	46.74	76.36	3.0
1977	12	22	14	57	00	46.86	76.94	3.5

LOWER ST. LAWRENCE (LSL)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1944	4	9	12	44	37	50.10	67.43	5.0
1944	6	23	6	37	52	49.42	67.75	5.0
1950	6	29	9	13	33	49.50	67.40	4.5
1951	9	19	8	19	37	49.30	66.25	4.5
1953	1	24	9	58	36	49.40	66.00	4.5
1953	9	14	22	52	57	49.40	65.30	4.5
1961	7	5	22	43	44	49.80	66.90	4.5
1964	7	1	21	41	30	49.43	67.42	3.8
1965	10	5	14	36	55	49.60	67.10	4.0
1966	1	14	15	29	25	48.90	67.70	4.0
1966	1	14	16	14	7	48.90	67.50	3.4
1966	7	12	1	6	38	49.50	66.00	3.3
1966	7	17	7	32	19	49.58	68.42	3.6
1966	7	24	22	19	46	49.63	68.55	3.7
1966	7	27	11	12	43	49.42	68.42	3.4
1966	8	20	13	13	33	49.58	68.33	3.5
1966	12	12	21	4	12	49.00	68.17	3.4
1967	9	30	22	39	51	49.30	65.90	4.5
1968	9	29	10	4	48	50.14	67.22	3.6
1972	8	22	19	17	49	49.60	66.40	4.0
1974	7	2	4	46	51	49.58	67.22	3.4
1974	12	27		50	12	49.14	67.44	3.5
1975	07	18	04	21	06	49.16	66.81	3.1
1975	10	21	20	50	02	49.13	68.13	3.1
1976	03	29	21	23	27	49.34	67.86	3.3
1976	05	15	21	06	52	49.84	68.62	3.3
1976	09	18	00	40	32	49.36	67.10	3.4
1977	08	08	23	06	12	49.70	67.08	3.8
1977	08	08	23	08	40	49.77	67.05	3.9
1977	08	08	23	29	27	49.70	67.02	2.8
1977	10	04	07	32	04	49.95	66.86	2.8

NORTHERN APPALACHIANS (NAP)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1937	9	30	7	58	10	45.47	65.83	4.5
1938	8	22	12	48	13	44.70	68.80	4.0
1940	3	28	11	42	34	44.70	69.90	3.8
1940	12	20	7	27	26	43.80	71.30	5.0
1940	12	24	13	43	44	43.80	71.30	5.0
1940	12	25	5	3	43	43.80	71.30	3.9
1940	12	27	19	56	9	43.80	71.30	3.9
1943	1	14	21	32	38	45.25	69.60	5.0
1943	3	14	14	2	27	43.70	71.57	3.9
1945	7	15	10	44	59	44.90	67.00	3.9
1947	12	28	19	58	18	45.27	69.25	4.0
1948	1	6	20	46	51	45.40	69.28	3.9
1949	10	5	2	33	47	44.80	70.50	4.0

NORTHERN APPALACHIANS (NAP)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1952	8	25		7		43.00	74.50	4.0
1953	3	31	12	58	34	44.07	73.12	3.9
1955	1	21	8	40		42.97	73.78	4.0
1957	4	23	19	42		44.40	72.00	4.0
1957	4	24		41	59	44.42	72.00	4.0
1957	4	26	11	40	6	43.60	70.40	4.0
1958	9	19	12	45		43.50	70.20	4.0
1961	1	29		49	39	46.38	66.93	3.8
1961	12	14	1	49	35	43.83	67.82	4.0
1962	4	10	14	30	48	44.15	73.05	4.5
1963	7	1	19	59	12	42.37	73.75	3.3
1963	12	4	21	32	34	43.60	71.60	3.7
1966	5	20		5	42	44.25	66.50	3.8
1966	7	24	1	59	58	44.50	67.60	3.6
1967	7	1	16	5	40	44.70	69.87	4.0
1968	5	27	19	21	56	46.90	66.66	3.3
1968	9	23	15	38	50	45.17	69.45	3.3
1970	8	8		10	30	45.80	66.12	3.3
1971	5	23	6	24	27	43.82	74.54	3.7
1971	5	23	9	29	59	43.94	74.55	3.6
1971	7	10	8	15	2	43.93	74.53	3.4
1973	6	15	1	9	5	45.39	71.03	5.0
1973	7	15	8	20	31	43.97	74.49	3.4
1973	7	16	8	41	58	43.76	74.47	3.3
1975	01	17	00	10	39	44.91	66.91	3.1
1975	02	28	18	40	21	46.39	66.01	2.9
1975	08	27	22	28	22	46.80	65.34	3.0
1975	10	15	03	26	17	45.11	65.89	3.1
1975	11	11	20	54	55	43.91	74.64	4.0
1976	03	08	18	08	40	46.78	64.96	2.8
1977	10	24	18	09	12	47.00	67.05	3.0
1979	04	18	02	34	14	43.95	69.75	4.0
1979	04	20	10	32	49	45.18	66.00	2.8
1980	09	08	05	59	55	44.68	69.00	3.2
1981	04	13	17	31	38	45.92	65.69	3.7
1981	05	08	18	56	13	45.90	65.97	2.8
1982	01	09	12	53	53	47.00	66.60	5.7
1982	01	09	13	09	40	47.00	66.60	3.4
1982	01	09	13	52	21	47.00	66.60	3.9
1982	01	09	16	36	45	47.00	66.60	5.1
1982	01	09	17	27	56	47.00	66.60	3.7
1982	01	09	17	37	37	47.00	66.60	3.3
1982	01	09	22	45	10	47.00	66.60	3.7
1982	01	11	21	41	09	47.00	66.60	5.5
1982	01	11	22	36	33	47.00	66.60	3.4
1982	01	12	01	58	01	47.00	66.60	3.5
1982	01	12	02	01	41	47.00	66.60	3.2
1982	01	12	11	49	31	47.00	66.60	2.9
1982	01	12	13	38	34	47.00	66.60	3.3
1982	01	13	17	56	46	47.00	66.60	4.0

NORTHERN APPALACHIANS (NAP)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1982	01	13	17	59	46	47.00	66.60	3.6
1982	01	15	12	37	39	47.00	66.60	3.7
1982	01	17	13	33	58	47.00	66.60	3.5
1982	01	23	08	56	49	47.00	66.60	2.8
1982	03	01	09	33	58	47.00	66.60	3.1
1982	03	04	06	06	33	47.00	66.60	2.8
1982	03	16	11	14	02	47.00	66.60	3.7
1982	03	20	03	08	11	47.00	66.60	3.1
1982	03	21	02	33	43	47.00	66.60	3.1
1982	03	26	05	36	41	47.00	66.60	2.8
1982	03	26	13	38	08	47.00	66.60	3.0
1982	03	31	21	02	22	47.00	66.60	4.8
1982	03	31	21	29	21	47.00	66.60	3.1
1982	04	02	13	50	13	47.00	66.60	4.3
1982	04	08	04	54	34	47.00	66.60	3.4
1982	04	10	01	59	00	47.00	66.60	2.9
1982	04	11	18	27	19	47.00	66.60	3.2
1982	04	18	22	47	21	47.00	66.60	4.1
1982	04	28	06	36	02	47.00	66.60	3.4
1982	05	02	01	42	45	47.00	66.60	3.1
1982	05	02	23	31	37	47.00	66.60	3.3
1982	05	06	16	28	08	47.00	66.60	4.0
1982	05	16	22	45	16	47.00	66.60	2.8
1982	05	28	06	24	26	47.00	66.60	3.0
1982	06	16	11	43	27	47.00	66.60	4.6
1982	06	16	16	15	41	47.00	66.60	3.0
1982	06	18	11	24	36	47.00	66.60	3.0
1982	06	25	06	47	10	47.00	66.60	2.9

LAURENTIAN SLOPE (LSP)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1929	11	18	20	32		44.50	56.30	7.0
1951	6	27	13	17	50	45.00	57.00	5.0
1954	8	28	15	23	1	45.17	56.87	5.2
1954	10	16	6	45	0	44.83	56.80	5.3
1965	11	28	23	26	10	45.57	57.90	4.2
1971	6	11	10	33	10	45.70	55.05	4.2
1975	3	31	17	8	2	44.70	56.22	4.6
1975	10	06	22	21	41	44.71	57.07	5.2
1977	7	16	11	37	43	44.52	55.83	3.8

ATTICA (ATT)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1929	8	12	11	24	48	42.87	78.35	5.5
1955	8	16	7	35		42.89	78.28	4.0
1965	7	16	11	6	55	43.20	78.50	3.5
1966	1	1	13	23	38	43.30	78.40	4.5
1967	6	13	19	8	54	43.30	78.00	4.5

EASTERN BACKGROUND (EBG)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1963	3	2	20	24	32	41.51	75.73	3.4
1963	5	19	19	14	18	43.50	75.23	3.5
1963	10	25	8	49	39	51.40	61.90	3.3
1964	1	20	18	57	55	46.83	71.33	3.9
1964	4	5	13	21	6	46.42	81.08	3.8
1964	6	27	19	17	46	47.75	79.17	3.7
1964	7	12			41	46.72	71.41	3.4
1964	8	12	9	35	27	50.47	64.87	3.7
1964	10	17	14	13	7	47.67	67.25	3.9
1965	1	8	12	29	45	48.00	78.50	3.5
1965	4	1	6	30	20	46.00	80.50	3.4
1965	10	18	12	10	17	53.13	79.27	3.6
1965	11	15	11	12	30	49.37	53.66	4.0
1965	12	19	0	49	17	51.17	80.83	4.5
1967	2	21	0	53	57	52.00	81.33	3.5
1967	2	22	14	21	55	50.50	63.33	3.5
1967	8	5	8	8	32	48.57	64.97	3.9
1967	9	17	1	19	38	50.67	75.25	4.0
1967	9	23	16	27	55	46.93	70.70	3.4
1967	10	25	7	5	31	50.15	63.52	3.4
1967	11	2	3	35	38	52.20	58.40	3.4
1968	10	10	20	10	41	45.80	81.66	3.4
1969	2	2	4	24	28	49.71	55.13	3.5
1969	8	5	21	53	23	47.66	52.29	3.4
1969	11	17	7	32	27	53.42	82.81	3.5
1971	08	15	06	17	15	47.46	49.53	4.6
1971	10	13	2	38	59	51.65	80.90	3.4
1975	9	2	6	21	17	48.29	69.74	3.3
1976	2	2	21	14	2	41.98	82.67	3.4
1976	08	28	19	23	30	50.10	48.85	4.0

BAFFIN BAY (BAB)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1933	11	20	23	21	32	73.00	70.75	7.3
1957	5	2	3	55	34	72.00	67.50	6.3
1969	11	27	8	25	24	73.50	70.64	4.2
1971	1	6	13	25	51	73.19	74.30	4.1
1971	6	2	20	47	49	74.92	67.56	4.6
1971	7	21	7	36	24	72.98	70.05	3.9
1972	2	3	6	21	52	73.00	71.68	4.3
1972	5	14	14	19	48	75.07	74.15	4.1
1972	5	30	19	47	26	71.66	64.91	3.8
1972	9	16	5	14	5	74.10	73.09	4.2
1974	2	2	20	11	26	74.19	72.45	4.4
1974	3	5	6	32	15	73.24	70.32	4.0
1974	3	8	17	48	8	74.12	69.16	4.8
1975	6	14	4	7	35	73.33	70.25	4.4
1976	3	20	0	47	29	73.15	69.98	4.7
1976	11	12	14	47	19	72.30	70.43	5.6

BAFFIN ISLAND (BAI)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1963	9	4	13	32	12	71.40	73.30	6.1
1966	12	26	4	12	58	71.50	74.67	4.9
1967	5	18	20	56	44	70.83	71.17	4.9
1968	1	22	3	1	39	70.97	73.22	4.3
1969	3	3	8	49	42	71.68	75.31	3.9
1969	3	3	9	15	8	71.66	75.27	3.8
1969	3	3	15	35	5	71.67	75.22	4.1
1969	3	4	1	52	9	71.65	75.88	4.4
1969	3	4	2	5	0	71.65	75.21	4.7
1969	3	6	16	37	53	71.69	75.08	3.9
1969	3	6	17	3	24	71.66	75.25	4.7
1969	3	6	17	8	3	71.74	75.34	4.2
1969	3	6	18	53	22	71.72	75.17	4.0
1969	3	10	11	35	51	71.66	75.13	3.8
1969	6	10	18	23	30	71.75	75.66	3.8
1970	10	20	17	0	48	70.96	73.24	4.3
1970	12	2	11	3	7	68.50	67.66	4.4
1971	1	16	23	11	16	71.68	75.24	4.2
1971	1	17	12	9	36	71.63	75.31	4.1
1971	1	17	15	48	7	71.62	75.34	3.9
1971	1	19	5	42	43	71.53	75.68	3.8
1971	3	22	8	58	5	68.40	68.32	3.9
1971	7	25	20	43	54	67.73	67.51	4.2
1971	7	31	22	57	52	71.76	76.43	4.0
1972	1	21	14	43	39	71.84	74.96	5.1
1972	5	10	20	1	41	71.60	74.96	4.0
1973	1	16	14	58	19	68.73	67.89	3.9
1973	4	24	3	55	5	71.38	71.22	4.7

BAFFIN ISLAND (BAI)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1973	5	29	9	3	55	71.74	74.92	4.4
1973	5	29	16	6	33	71.71	75.26	4.5
1973	5	29	16	7	0	71.70	75.30	4.0
1973	8	17	5	31	26	71.38	70.78	4.1
1973	12	7	8	2	49	69.02	69.81	4.1
1973	12	7	8	3	31	69.05	69.93	4.3
1973	12	14	18	21	32	71.70	75.24	4.2
1974	7	19	16	38	23	71.97	75.56	4.0
1975	5	3	16	19	32	71.14	73.20	4.0
1975	6	30	18	48	55	71.44	71.19	5.2
1976	8	17	4	30	2	69.02	70.27	3.8
1976	10	11	20	43	2	68.62	67.82	3.8
1977	12	19	07	42	06	71.80	75.08	4.0

LABRADOR SEA (LAB)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1969	9	27	22	53	58	56.52	57.49	4.1
1969	11	24	21	14	14	60.60	58.80	5.0
1969	11	30	14	38	6	60.55	59.22	4.2
1970	7	3	0	32	36	60.89	60.47	4.2
1971	1	12	17	36	4	62.31	62.33	3.9
1971	4	16	1	31	45	61.75	60.68	4.3
1971	7	13	1	32	12	60.63	57.45	3.8
1971	12	7	12	4	18	55.09	54.51	5.6
1972	1	25	2	40	1	55.14	54.42	4.5
1973	8	27	1	49	36	60.07	57.91	4.4
1973	10	12	3	54	28	61.34	59.99	4.4
1975	12	13	9	24	27	57.94	52.25	4.5
1976	5	26	18	26	33	55.47	52.74	4.4
1977	9	24	17	19	44	58.25	54.24	4.8
1977	11	5	8	49	31	59.05	60.61	4.2

EASTERN ARCTIC BACKGROUND (EAB)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1951	4	22	12	36	16	76.00	73.00	5.7
1957	7	21	8	53	31	68.90	59.40	5.7
1971	5	31	4	58	52	68.92	53.88	4.2
1971	7	4	5	32	50	68.91	53.09	3.9
1972	4	15	14	29	49	72.48	55.37	4.0
1972	5	28	23	8	3	76.16	72.30	4.2
1973	1	20	0	24	17	68.53	59.63	4.9
1975	7	20	6	28	38	75.23	62.39	4.4
1976	4	4	3	4	26	68.78	54.93	4.0
1976	11	2	13	30	2	69.77	53.70	4.4

EASTERN ARCTIC BACKGROUND (EAB) (CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1977	06	01	05	27	33	69.20	53.91	3.8

GUSTAF-LOUGHEED ARCH (GLA)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1970	5	25	7	24	10	76.75	105.77	4.3
1972	7	7	10	9	0	79.86	107.36	4.4
1972	7	21	21	11	35	79.97	108.45	4.0
1972	9	30	22	51	30	79.88	107.72	5.0
1972	11	17	10	36	15	76.69	106.44	4.8
1972	11	17	10	53	16	76.71	106.32	4.9
1972	11	19	17	33	44	76.55	106.33	5.6
1972	11	19	18	45	48	76.47	106.38	4.7
1972	11	20	6	7	13	76.57	106.02	3.9
1972	11	21	10	6	27	76.58	106.02	5.7
1972	11	21	13	58	18	76.57	106.29	4.1
1972	11	21	17	42	49	76.54	106.46	4.2
1972	11	25	19	43	37	76.55	106.44	3.9
1972	12	7	21	48	22	76.64	106.46	3.9
1972	12	13	18	38	39	76.58	106.63	4.7
1972	12	13	19	55	19	76.54	106.71	4.0
1972	12	20	17	30	21	76.56	106.75	4.7
1972	12	22	19	32	3	76.52	106.51	3.9
1972	12	27	22	59	26	76.80	106.49	5.4
1972	12	28	12	18	44	76.76	106.47	4.1
1972	12	28	13	49	43	76.79	106.64	3.9
1972	12	28	14	1	2	76.73	106.61	4.1
1972	12	28	14	36	5	76.80	106.16	5.1
1972	12	29	20	1	44	76.77	106.35	4.0
1973	1	1	0	21	2	76.68	106.22	4.0
1973	1	3	8	32	3	76.69	105.90	3.8
1973	1	9	13	17	41	76.62	105.90	4.4
1973	1	9	13	23	20	76.61	106.22	4.4
1973	1	31	21	33	23	76.74	106.39	4.4
1973	2	12	10	56	33	76.42	107.03	3.8
1973	2	14	3	38	38	76.77	106.14	4.0
1973	3	5	5	20	59	76.54	106.76	3.8
1973	3	16	11	13	37	76.50	106.42	3.8
1973	4	30	21	27	24	76.59	106.29	3.8
1973	5	3	5	42	32	76.48	106.41	4.9
1973	5	15	15	40	52	76.58	106.61	3.9
1973	8	19	10	36	42	76.52	106.45	3.8
1974	2	21	15	30	50	76.70	106.54	4.2
1974	3	28	17	30	17	76.68	106.28	4.6
1974	9	14	2	2	56	80.03	107.89	4.6
1974	9	24	9	13	59	76.73	106.37	3.9
1974	9	24	10	23	54	76.85	106.36	3.9
1974	11	8	11	28	13	79.73	108.92	4.1

GUSTAF-LOUGHEED ARCH (GLA)

(CONTINUED)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1974	12	27	22	53	44	76.62	106.06	4.9
1974	12	27	23	0	16	76.68	106.38	4.8
1974	12	27	23	21	38	76.65	106.30	4.6
1974	12	28	6	17	46	76.70	106.19	4.3
1975	1	3	2	3	50	76.63	106.26	4.3
1975	1	3	11	8	52	76.71	106.44	4.7
1976	3	14	14	4	10	79.92	107.77	4.4
1976	8	27	8	18	23	76.65	106.11	3.8
1976	9	16	10	14	34	76.64	106.55	4.6
1977	02	18	19	24	09	76.69	106.32	3.9
1977	05	25	23	01	04	77.38	104.56	4.2
1978	02	05	16	07	12	78.24	107.33	4.8
1978	02	05	16	13	40	78.37	107.45	4.2
1978	02	19	22	59	12	78.25	107.42	4.0
1978	02	23	23	32	38	78.32	107.76	3.8
1978	02	26	20	50	40	78.28	107.62	4.1
1978	03	08	21	15	26	78.26	107.57	4.3
1978	04	16	03	27	41	78.35	107.53	4.1

SVERDRUP (SVD)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1968	11	11	19	45	2	79.03	95.12	3.8
1969	4	23	18	59	42	75.64	102.22	3.9
1970	12	24	12	19	48	75.61	101.11	4.0
1971	2	3	13	47	39	77.17	118.18	3.8
1974	9	7	23	50	39	76.74	100.88	3.8
1975	3	8	5	20	34	79.82	94.07	5.2
1975	3	28	4	18	8	78.09	110.08	3.8
1975	4	15	12	17	40	81.19	87.14	4.1
1975	4	23	18	31	9	79.73	94.01	4.0
1975	5	22	21	54	41	76.09	121.91	3.8
1975	6	3	2	38	56	76.22	119.44	4.0
1975	7	15	21	10	21	76.42	120.11	3.8
1975	9	19	11	34	2	80.03	93.82	3.8
1976	2	12	3	29	16	82.14	76.38	4.2
1976	3	15	8	50	30	81.28	84.68	3.8
1977	07	08	05	27	31	75.61	105.74	4.0

BOOTHIA-UNGAVA (BOU)

YR	MO	DA	HR	MN	SC	LAT	LONG	M
1959	1	30	5	17	32	61.00	78.50	5.9
1960	9	6	21	24	26	64.70	86.40	5.5
1966	3	22	22	10	3	64.75	88.00	5.1
1968	12	19	16	49	18	67.47	91.41	3.8
1971	3	17	16	47	29	74.82	94.37	3.9
1971	6	27	3	34	23	73.27	95.77	4.0
1971	7	24	22	25	55	73.65	96.13	4.0
1971	9	20	23	1	41	73.81	92.52	3.9
1971	9	26	7	1	6	59.95	73.67	3.9
1971	10	2	3	19	28	64.20	86.67	5.1
1972	1	22	12	21	25	72.44	93.68	4.2
1972	1	24	18	26	37	64.60	88.13	4.0
1972	5	18	5	8	47	74.19	95.77	4.5
1972	7	4	10	12	21	73.73	96.65	4.0
1972	10	27	1	59	20	67.05	94.55	4.4
1973	1	25	14	52	3	65.60	89.07	3.8
1974	4	21	4	48	50	74.33	93.91	4.9
1975	5	22	15	5	52	67.23	92.14	4.9
1976	2	29	16	18	42	60.33	76.59	3.9
1976	4	14	17	16	12	64.33	89.93	4.3
1976	8	6	11	14	16	63.03	86.78	3.9
1976	12	6	6	9	52	65.22	87.88	4.0

APPENDIX B

Listing of Modified McGuire (1976) Seismic Risk Program

```

PROGRAM CANRSK(INPUT,OUTPUT,TAPE1=INPUT,TAPE6=OUTPUT,TAPE2)
COMMON NRD,NWR,RSKTI(12)
COMMON C1,C2,C3,SIG,RZERO,RONE,AAA,BBB
COMMON NGS,NRS(23),AMO(24),AM1(24),LORS(24)
COMMON BETA(24),RATE(24),COEF(24),FDEPTH(24)
COMMON NSTEPO,NSTEP1
COMMON INDIC(4),AREA(23,11),X(23,11,2),Y(23,11,2)
COMMON/DEBG/SXNOT,SYNOT,ERRBND,NSTEPMX,LERR,RZ2,LERR2,RISKER
COMMON /MDATA/NSTEP,JPRNT,JPRNT2,JPRNT3,NLEI,TI(12),RISKS(9),
1ATTEN(8,2,2)
COMMON/LERRS/INCLUD,INCLD2,LNLEI,LRISKS,RKRATO
COMMON/CNM/RNAME(6),SNAME(6),TIFS(8,2),RZ2S(23),JEW,NRG,A,B
1,IN,JN,NRGL
DIMENSION CLAT(7)
LOGICAL LERR,INCLUD,LERR2,INCLD2,LNLEI,LRISKS,LPR
DATA INCLUD,INCLD2,LNLEI,LRISKS,ERRBND,NSTEPMX,RKRATO,NSTEP
1/.F.,.F.,.F.,.F.,20.,40,0.05,10/
DATA NLEI,TI/10,-2.,-1.,0.,1.,2.,3.,4.,5.,6.,7.,0.,0./
DATA RISKS/0.01,0.005,0.002105,0.001,0.,0.,0.,0.,0./
DATA ATTEN/-1.0788,1.3,-1.1,0.7,0.,0.,1000.,0.0001,
1      -8.623,2.3,-1.0,0.7,0.,0.,1000.,0.0001,
2      0.,1.3,-1.5,0.7,0.,0.,1000.,0.0001,
3      -7.824,2.3,-1.3,0.7,0.,0.,1000.,0.0001/
DATA CLAT/-70.,-70.,-80.,-105.,-123.,-123.,-138./
IN=1$NWR=6$NRD=2$JN=6
NRGL=0
WRITE(JN,1000)
1000 FORMAT(" CANADIAN SEISMIC RISK PROGRAM RUNNING"/)
9 WRITE(JN,50)
50 FORMAT(" NAME OF REQUESTOR ?",15H: "END" TO STOP/)
LPR=.TRUE.
READ(IN,1)RNAME
1 FORMAT(6A10)
IF(EOF(IN))99,2
2 IF(RNAME(1).EQ.10HEND ) GO TO 99
IF(RNAME(1).EQ.10H ) GO TO 9
IF(RNAME(1).EQ.10H/ )GO TO 60
5 WRITE(JN,51)
51 FORMAT(" NAME OF SITE ?"/)
READ(IN,1)SNAME
IF(SNAME(1).EQ.10H )GO TO 5
6 WRITE(JN,53)
53 FORMAT(" LAT & LONG ?"/)
READ *,A,B
IF(A.LT.0.)GO TO 9$IF(A.EQ.0..OR.B.EQ.0.)GO TO 6
IF(.NOT.LPR) GO TO 21
WRITE(JN,7)A,B
7 FORMAT(1X,F7.3,"N ",F8.3,"W ? Y/N")
READ(IN,8)YN
8 FORMAT(A1)
IF(YN.EQ.1HY) GO TO 21
GO TO 6
60 LPR=.FALSE.$GO TO 6
21 A=ABS(A)$B=ABS(B)
IF(B.LT.49..OR.B.GT.145.)GO TO 30

```

```

IF(B.LE.145..AND.B.GE.91..AND.A.GE.68..AND.A.LE.82.)GO TO 10
IF(B.GE.110.)GO TO 20
JEW=1
IF(A.LT.41..OR.A.GT.85.)GO TO 30
IF(B.LE.86..AND.A.GE.41..AND.A.LT.50.) GO TO 11
IF(B.LE.90..AND.A.GE.50..AND.A.LT.60.) GO TO 12
IF(B.LE.105..AND.A.GE.60..AND.A.LE.85.) GO TO 13
IF(A.GE.46..AND.A.LE.51..AND.B.GE.100.) GO TO 14
GO TO 40
20 JEW=2
IF(A.LT.47.)GO TO 30
IF(A.GE.47..AND.A.LT.53.) GO TO 15
IF(A.GE.53..AND.A.LT.60.) GO TO 16
IF(A.GE.60..AND.A.LE.68.) GO TO 17
GO TO 40
30 WRITE(JN,31)A,B
31 FORMAT("/" SITE AT",F6.2," LAT  ",F7.2," LONG"/
1" IS OUTSIDE AREA WHERE RISK CAN BE DEFINED"/)
GO TO 9
10 NRG=8$CALL INLCCM(-115.)$CALL NWNERK$GO TO 22
40 WRITE(JN,33)A,B
33 FORMAT(" THE RISK AT SITE",F6.2," LAT  ",F7.2," LONG"/
1" IS INSIGNIFICANT"/)
GO TO 9
11 NRG=1$GO TO 23
12 NRG=2$GO TO 23
13 NRG=3$GO TO 23
14 NRG=4$GO TO 23
15 NRG=5$GO TO 23
16 NRG=6$GO TO 23
17 NRG=7
23 IF(NRG.EQ.NRGL)GO TO 24$CALL INLCCM(CLAT(NRG))
24 CALL SRISK
22 IF(LPR)CALL OUTFRM$IF(.NOT.LPR) CALL SHOUT$IF(LPR)GO TO 9$GO TO 6
99 STOP$END

```



```

SUBROUTINE AREAS(X1,Y1,X2,Y2,X3,Y3,X4,Y4,AREA)
C   SUBROUTINE TO CALCULATE AREA OF ARBITRARY QUADRILATERAL,
C   WHERE (X1,Y1) AND (X4,Y4) ARE OPPOSITE CORNERS.
C   LOCATE INTERSECTIONS OF DIAGONALS:
IF (X4-X1-0.001) 20,30,30
20 IF (X4-X1+0.001) 30,30,25
25 XX=X1
A2=(Y3-Y2)/(X3-X2)
YY=(XX-X3)*A2 +Y3
DIST1=Y4-Y1
IF (DIST1) 26,27,27
26 DIST1=-DIST1
27 DIST2=SQRT((X3-X2)*(X3-X2) + (Y3-Y2)*(Y3-Y2))
GO TO 100
30 IF (X3-X2-0.001) 50,70,70
50 IF (X3-X2+0.001) 70,70,65
65 XX=X3
A1=(Y4-Y1)/(X4-X1)
YY=(XX-X4)*A1+Y4
DIST2=Y3-Y2
IF (DIST2) 66,67,67
66 DIST2=-DIST2
67 DIST1=SQRT((X4-X1)*(X4-X1) + (Y4-Y1)*(Y4-Y1))
GO TO 100
70 A1=(Y4-Y1)/(X4-X1)
A2=(Y3-Y2)/(X3-X2)
XX=(Y2-Y1+A1*X1-A2*X2)/(A1-A2)
YY=A1*(XX-X1) + Y1
DIST1=SQRT((X4-X1)*(X4-X1) + (Y4-Y1)*(Y4-Y1))
DIST2=SQRT((X3-X2)*(X3-X2) + (Y3-Y2)*(Y3-Y2))
C   CALCULATE LENGTH OF SIDES OF SUB-TRIANGLE
100 SIDE1=SQRT((XX-X1)*(XX-X1) + (YY-Y1)*(YY-Y1))
SIDE2=SQRT((XX-X2)*(XX-X2) + (YY-Y2)*(YY-Y2))
SIDE3=SQRT((X1-X2)*(X1-X2) + (Y1-Y2)*(Y1-Y2))
C   SOLUTION ACCORDING TO C.R.C. HANDBOOK UNDER
C   'MENSURATION FORMULAE' AND 'TRIGONOMETRIC FORMULAE'
SS=(SIDE1+SIDE2+SIDE3)/2.
SINANG=2.*SQRT(SS*(SS-SIDE1)*(SS-SIDE2)*(SS-SIDE3))/(SIDE1*SIDE2)
AREA=0.5*DIST1*DIST2*SINANG
RETURN
END
SUBROUTINE BETWEEN(X1,Y1,X2,Y2,XP,YP,INDIC,IANS)
C   SUBROUTINE TO DETERMINE IF (XP,YP) LIES BETWEEN
C   (X1,Y1) AND (X2,Y2).
IF(INDIC.LT.1.OR.INDIC.GT.2) GO TO 300
IF(INDIC.EQ.2) GO TO 200
IF (X1-XP) 110,410,120
110 IF (X2-XP) 420,420,410
120 IF (X2-XP) 410,420,420
200 IF (Y1-YP) 210,410,220
210 IF (Y2-YP) 420,420,410
220 IF (Y2-YP) 410,420,420
C   ERROR RETURN
300 IANS=0
GO TO 500

```

```

C      (XP,YP) LIES BETWEEN END POINTS, I.E. ON SOURCE BOUNDARY
410  IANS=1
      GO TO 500
C      (XP,YP) DOESN'T LIE BETWEEN END POINTS, I.E. IT'S OUTSIDE SOURC
420  IANS=-1
500  RETURN
      END
      SUBROUTINE CIRCLE(RC,INGS,FRAREA,RSK)
      COMMON NRD,NWR,RSKTI(12)
      COMMON C1,C2,C3,SIG,RZERO,RONE,AAA,BBB
      COMMON NGS,NRS(23),AMO(24),AM1(24),LORS(24)
      COMMON BETA(24),RATE(24),COEF(24),FDEPTH(24)
      COMMON NSTEP0,NSTEP1
      COMMON INDIC(4),AREA(23,11)
      COMMON /MDATA/NSTEP,JPRNT,JPRNT2,JPRNT3,NLEI,TI(12),RISKS(9),
1ATTEN(8,2,2)
      DIMENSION RSK(12)
C      SUBROUTINE TO CALCULATE RISK FROM A CIRCULAR
C      SOURCE WITH CENTER AT SITE, RADIUS RC.
      NRC=RC
C      CHOOSE STEP SIZE:
C      STEP SIZE = NSTEPI UNLESS RESULTING STEP SIZE IS
C      LESS THAN ONE KILOMETRE, IN WHICH CASE RC+1
C      STEPS ARE USED.
      IF (NRC-NSTEP1) 10,12,12
10   NSTEPX=NRC+1
      GO TO 14
12   NSTEPX=NSTEP1
14   ANSTEP=NSTEPX
      DO 90 II=1,NSTEPX
      AI=II
      R=((AI-0.5)*RC)/ANSTEP
      ANAREA=6.2831853072 *R*RC/ANSTEP
      RATEI=RATE(INGS)*ANAREA*FRAREA
      DO 80 JJ=1,NLEI
      CALL RISK1(TI(JJ),R,INGS,RISK)
      RSK(JJ)=RSK(JJ)+RISK*RATEI
80   CONTINUE
90   CONTINUE
      RETURN
      END
      SUBROUTINE COMBINI
      COMMON NRD,NWR,RSKTI(12)
      COMMON C1,C2,C3,SIG,RZERO,RONE,AAA,BBB
      COMMON NGS,NRS(23),AMO(24),AM1(24),LORS(24)
      COMMON BETA(24),RATE(24),COEF(24),FDEPTH(24)
      COMMON NSTEP0,NSTEP1
      COMMON INDIC(4),AREA(23,11),X(23,11,2),Y(23,11,2)
      COMMON/DEBG/SXNOT,SYNOT,ERRBND,NSTEPX,LERR,RZ2,LERR2,RISKER
      COMMON /MDATA/NSTEP,JPRNT,JPRNT2,JPRNT3,NLEI,TI(12),RISKS(9),
1ATTEN(8,2,2)
      COMMON/SRSKC/SRSK(12,4)
      COMMON/CNM/RNAME(6),SNAME(6),TIFS(8,2),RZ2S(23),JEW,NRG,SLAT,SLONG
1,IN,JN,NRGL
      DIMENSION TIF(8)

```

```

LOGICAL LERR,LERR2
DO 424 J=1,2
DO 13 I=1,NLEI
13 RSKTI(I)=SRSK(I,J)+SRSK(I,J+2)
DO 620 I=1,NLEI
620 RSKTI(I)=1.-EXP(-RSKTI(I))
C     ESTIMATE INTENSITIES AT RISKS DESIRED.
RISKS(9)=0.0
IA=0
IF(RISKS(1)-0.0000000001) 700,700,625
625 DO 630 IRK=1,8
IF (RISKS(IRK)-RSKTI(1)) 640,640,630
630 TIF(IRK)=1000000.
GO TO 700
640 IA=IA+1
IF (IA-NLEI) 650,645,645
645 TIF(IRK)=1000000.
IRK=IRK+1
IF (RISKS(IRK)-0.0000000001)680,680,645
650 IF(RISKS(IRK)-RSKTI(IA+1))640,655,655
655 TIF(IRK)=(ALOG(RSKTI(IA)/RISKS(IRK)))
1 /((ALOG(RSKTI(IA)/RSKTI(IA+1)))
TIF(IRK)=TIF(IA)+TIF(IRK)*(TIF(IA+1)-TIF(IA))
IRK=IRK+1
IF (RISKS(IRK)-0.0000000001)680,680,660
660 IF(RISKS(IRK)-RSKTI(IA+1)) 640,655,655
680 IRK=IRK-1
DO 685 I=1,IRK
IF (TIF(I)-999999.)683,685,685
683 TIF(I)=EXP(TIF(I))
685 CONTINUE
DO 105 I=1,IRK
105 TIFS(I,J)=TIF(I)
700 CONTINUE
424 CONTINUE
RETURN$END
SUBROUTINE ERISK(AMZ,AMM,C1,C2,C3,RLN,SIG,BETA,TIC,
1 G1,G2,G3,G4,CON1,CON2,CON3)
C     EVALUATE RISK ASSOCIATED WITH EXPONENTIAL MAGNITUDE LAW
C     FOR MAGNITUDES BETWEEN AMZ AND AMM.
Z=(TIC-C1-C2*AMM-C3*RLN)/SIG
CALL NDTR(Z,G1,D)
Z=Z-BETA*SIG/C2
CALL NDTR(Z,G3,D)
Z=(TIC-C1-C2*AMZ-C3*RLN)/SIG
CALL NDTR(Z,G2,D)
Z=Z-BETA*SIG/C2
CALL NDTR(Z,G4,D)
IF (C2-0.001) 10,10,20
10 CON1=1000000000.
CON3=CON1
GO TO 30
20 CON1=((BETA*BETA*SIG*SIG)/(2.*C2*C2))+(BETA*AMZ)
1 +((C1-TIC)*BETA/C2)
CON3=CON1+BETA*(AMM-AMZ)

```



```

CON1=EXP(CON1)
CON3=EXP(CON3)
30 CON2=BETA*C3/C2
R=EXP(RLN)
CON2=R**CON2
RETURN
END
SUBROUTINE INLCCM(SLONG)
C SLONG CAN. CENTRAL LONGITUDE
COMMON/CTAS/SCALEK,CONE,TLONG,RO,RADCON
DATA RAD,E,E2,A,ED2/0.01745329252,12.15482511000,0.00676865800218,
C 637820640.1,0.041135927122/
DATA TSPAR,TNPAR,TLAT,TSCALE
1 / 49.,77., 63., 1E5/
C PARAMS 1,2 +3 ARE STANDARD LATs. LONGs. FOR CAN. PROJECTION
C 1E5 IS TO CONVERT FROM AN EASTING(X) AND NORTHING(Y) TO KM.
C REGCLN REMAINS ONLY INIT-PARAM. FIXING CENTRAL MERIDIAN, I.E. VERT
C IN PLOTTING ROUTINE, WHERE X=C
C INITIALIZATION OF LAMBERT CONFORMAL CONIC PROJECTION
P=TSPAR*RAD
SP=SIN(P)
ANSSPS=COS(P)/((1.+SP)*((E+SP)/(E-SP))**ED2)
P=TNPAR*RAD
SP=SIN(P)
ANSSPN=COS(P)/((1.+SP)*((E+SP)/(E-SP))**ED2)
SPS=TSPAR*RAD
SPN=TNPAR*RAD
TLONG=SLONG
SS=SIN(SPS)
QN1=SQRT(1.-SS*SS*E2)
SS=SIN(SPN)
QN2=SQRT(1.-SS*SS*E2)
CONEN=ALOG(COS(SPS)*QN2/(COS(SPN)*QN1))
CONE=CONEN/ALOG(ANSSPS/ANSSPN)
RADCON=RAD*CONE
SCALEK=A*COS(SPS)/(CONE*ANSSPS**CONE*TSCALE*QN1)
P=TLAT*RAD
SP=SIN(P)
RO=SCALEK*(COS(P)/((1.+SP)*((E+SP)/(E-SP))**ED2)**CONE)
RETURN
END
SUBROUTINE INSIDE(XNOT,YNOT,INGS,INSS,XL,YL,XR,YR,AA,BB)
COMMON NRD,NWR,RSKTI(12)
COMMON C1,C2,C3,SIG,RZERO,RONE,AAA,BBB
COMMON NGS,NRS(23),AMO(24),AM1(24),LORS(24)
COMMON BETA(24),RATE(24),COEF(24),FDEPTH(24)
COMMON NSTEPO,NSTEP1
COMMON INDIC(4),AREA(23,11)
COMMON/DEBG/SXNOT,SYNOT,ERRBND,NSTEPMX,LERR,RZ2,LERR2,RISKER
COMMON /MDATA/NSTEP,JPRNT,JPRNT2,JPRNT3,NLEI,TI(12),RISKS(9),
1ATTEN(8,2,2)
DIMENSION XL(4),YL(4),XR(4),YR(4),AA(4),BB(4),XC(4),YC(4)
DIMENSION RSK(10)
LOGICAL LERR,LERR2
NSTEPI=NSTEP1

```

```

C          SUBROUTINE FOR CALCULATING RISK WHEN SITE IS INSIDE
C          SOURCE AREA.
C          REFER TO DOCUMENTATION FOR ALGORITHM USED TO CHOOSE
C          NUMBER OF INTEGRATION STEPS.
      3 DO 50 II=1,NLEI
50    RSK(II)=0.
      APPROX=0.
      RC2=100000000000.
      RF2=0.
C          FIND CLOSEST SIDE AND FARTHEST POINT
      DO 160 II=1,4
      IF(INDIC(II).EQ.1) GO TO 120
      XS=XL(II)
      YS=YNOT
      GO TO 140
C          IS SLOPE ZERO?
120    IF (AA(II)-0.001) 121,125,125
121    IF (AA(II)+0.001) 125,125,122
122    XS=XNOT
      YS=YL(II)
      GO TO 140
C          SLOPE NOT ZERO
125    XS=(YNOT+(XNOT/AA(II))-BB(II))/(AA(II)+1./AA(II))
      YS=((XNOT-XS)/AA(II))+YNOT
C          CALCULATE SQUARE OF DISTANCE BETWEEN SITE AND CLOSEST POINT.
140    DIST=(XNOT-XS)*(XNOT-XS)+(YNOT-YS)*(YNOT-YS)
      IF (DIST-RC2) 151,152,152
151    RC=SQRT(DIST)
      RC2=DIST
      ICLO=II
C          CALCULATE DISTANCE BETWEEN SITE AND LEFT HAND POINT ON SIDE
152    DIST=(XNOT-XL(II))*(XNOT-XL(II))+(YNOT-YL(II))*(YNOT-YL(II))
      IF (RF2-DIST) 154,160,160
154    RF=SQRT(DIST)
      RF2=DIST
      IFAR=II
160    CONTINUE
C          DETERMINE AZIMUTH OF FARTHEST POINT WITH RESPECT TO SITE
      AZIMF=ACOS((XL(IFAR)-XNOT)/RF)
      IF (YL(IFAR)-YNOT) 162,164,164
162    AZIMF=6.2831853072 - AZIMF
164    CONTINUE
C          RC IS NOW DISTANCE FROM SITE TO CLOSEST SIDE.
C          RF IS NOW DISTANCE FROM SITE TO FARTHEST CORNER.
      IF (RC-0.01) 200,200,170
C          CALL SUBROUTINE CIRCLE TO CALCULATE RISK FROM CIRCULAR
C          SOURCE WITH RADIUS RC
170    NTOT=NRS(INGS)+1
      FRAREA=AREA(INGS,INSS)/AREA(INGS,NTOT)
      CALL CIRCLE(RC,INGS,FRAREA,RSK)
      APPROX=3.1415926536*RC*RC
C          LOOP ON R TO CALCULATE RISK FROM RC TO RF
200    AN=NSTEP1
C          PICK STEP SIZE BASED ON FRACTION OF AREA LEFT
      FRLEFT=(AREA(INGS,INSS)-APPROX)/AREA(INGS,INSS)

```

```

NSTEPX=FRLEFT*AN + 1.
AN=NSTEPX
STSIZE=(RF-RC)/AN
DO 500 ISTEP=1,NSTEPX
AI=ISTEP
R=RC+(AI-0.5)*STSIZE
NPT=0
ANGLE=0.
C      LOOP ON EACH SIDE
DO 400 II=1,4
IF(INDIC(II).EQ.1) GO TO 330
C      SIDE II IS VERTICAL, DOES CIRCLE (RADIUS R) INTERSECT IT?
A=XL(II)-XNOT
IF (A) 322,322,323
322 IF (R+A) 400,400,324
323 IF (R-A) 400,400,324
C      COMPUTE 2 INTERSECTION POINTS
324 X1=XL(II)
B=SQRT(R*R-(X1-XNOT)*(X1-XNOT))
Y1=YNOT+B
X2=XL(II)
Y2=YNOT-B
GO TO 341
330 A=1.+AA(II)*AA(II)
B=2.*(-XNOT+AA(II)*(BB(II)-YNOT))
C=XNOT*XNOT+YNOT*YNOT+BB(II)*(BB(II)-2.*YNOT)-R*R
D=B*B-4.*A*C
IF (D) 400,400,340
C      THERE ARE 2 INTERSECTION, CALCULATE THEIR COORDINATES.
340 D=SQRT(D)
X1=(-B+D)/(2.*A)
Y1=AA(II)*X1+BB(II)
X2=(-B-D)/(2.*A)
Y2=AA(II)*X2+BB(II)
C      SEE IF (X1,Y1) IS ON BOUNDARY
341 CALL BETWEN(XL(II),YL(II),XR(II),YR(II),X1,Y1,INDIC(II),IANS)
IF (IANS) 350,342,345
342 NERROR=4
GO TO 800
C      IS SECOND POINT ALSO ON BOUNDARY?
345 CALL BETWEN(XL(II),YL(II),XR(II),YR(II),X2,Y2,INDIC(II),IANS)
IF (IANS) 348,346,360
346 NERROR=5
GO TO 800
C      STORE FIRST POINT ONLY
348 NPT=NPT+1
XC(NPT)=X1
YC(NPT)=Y1
GO TO 400
C      SEE IF SECOND POINT ONLY IS ON BOUNDARY
350 CALL BETWEN(XL(II),YL(II),XR(II),YR(II),X2,Y2,INDIC(II),IANS)
IF (IANS) 400,352,354
352 NERROR=6
GO TO 800
354 NPT=NPT+1

```



```

XC(NPT)=X2
YC(NPT)=Y2
GO TO 400
C      TWO INTERSECTION POINTS ON ONE SIDE BOTH LIE ON BOUNDARY,
C      CALCULATE ANGLE BETWEEN THEM.
360  CONTINUE
380  AD=SQRT((X1-X2)*(X1-X2)+(Y1-Y2)*(Y1-Y2))
    IF (AD.GT.ABS(2.*R)) AD=2.*ABS(R)
    ANGLE=2.*ASIN(AD/(2.*R))+ANGLE
400  CONTINUE
    IF (NPT) 402,404,408
402  NERROR=7
    GO TO 800
404  IF (ANGLE-0.001) 402,406,406
C      FOLLOWING IS FOR CASE OF NO SINGLE INTERSECTION POINTS;
C      ANGLE IS 2 * PI - ANGLE CALCULATED SO FAR.
406  PANGLE=6.2831853072-ANGLE
    GO TO 460
408  IF(NPT.LT.1.OR.NPT.GT.4) GO TO 409
    GO TO (402,410,402,440),NPT
409  NERROR=8
    GO TO 800
C      2 INTERSECTION POINTS; DETERMINE AZIMUTHS.
410  IF (XC(1)-XNOT-R) 414,413,411
411  IF (XC(1)-XNOT-R-0.001) 413,413,412
412  NERROR=18
    GO TO 800
413  AZIM1=0.0
    GO TO 418
414  IF (XC(1)-XNOT+R) 415,416,417
415  IF (XC(1)-XNOT+R+0.001) 412,416,416
416  AZIM1=3.1415926536
    GO TO 420
417  AZIM1=ACOS((XC(1)-XNOT)/R)
418  IF (YC(1)-YNOT) 419,420,420
419  AZIM1=6.2831853072 - AZIM1
420  IF (XC(2)-XNOT-R) 424,423,421
421  IF (XC(2)-XNOT-R-0.001) 423,423,422
422  NERROR=19
    GO TO 800
423  AZIM2=0.0
    GO TO 428
424  IF (XC(2)-XNOT+R) 425,426,427
425  IF (XC(2)-XNOT+R+0.001) 422,426,426
426  AZIM2=3.1415926536
    GO TO 430
427  AZIM2=ACOS((XC(2)-XNOT)/R)
428  IF (YC(2)-YNOT) 429,430,430
429  AZIM2=6.2831853072 -AZIM2
430  PANGLE=AZIM2-AZIM1
    IF (PANGLE) 431,439,435
431  IF (AZIM1-AZIM2) 432,439,433
432  PANGLE=6.2831853072 +PANGLE -ANGLE
    GO TO 460
433  IF (AZIM1-AZIM2) 432,439,434

```

```

434 PANGLE=-PANGLE-ANGLE
GO TO 460
435 IF (AZIM2-AZIMF) 436,439,437
436 PANGLE=6.2831853072 -PANGLE -ANGLE
GO TO 460
437 IF (AZIMF-AZIM1) 436,439,438
438 PANGLE=PANGLE-ANGLE
GO TO 460
439 NERROR=9
GO TO 800
C      FOUR INTERSECTION POINTS (EACH ON A DIFFERENT SIDE).
C      DETERMINE ANGLE BY FINDING CLOSEST 2 INTERSECTIONS TO
C      FARTHEST CORNER, CALCULATE ANGLE BETWEEN, AND ADD ANGLE
C      BETWEEN OTHER TWO INTERSECTIONS.
440 DIST1=10000000000.
I1=0
I2=0
I3=0
I4=0
DO 450 JJ=1,4
DIST=(XL(IFAR)-XC(JJ))*(XL(IFAR)-XC(JJ))
1  +(YL(IFAR)-YC(JJ))*(YL(IFAR)-YC(JJ))
IF (DIST-DIST1) 442,444,444
442 DIST2=DIST1
DIST1=DIST
I4=I3
I3=I2
I2=I1
I1=JJ
GO TO 450
444 IF (DIST-DIST2) 445,446,446
445 DIST2=DIST
I4=I3
I3=I2
I2=JJ
GO TO 450
446 I4=I3
I3=JJ
450 CONTINUE
C      CALCULATE ANGLE BETWEEN 2 CLOSEST POINTS TO FARTHEST CORNER
AD=SQRT((XC(I1)-XC(I2))*(XC(I1)-XC(I2))
1  + (YC(I1)-YC(I2))*(YC(I1)-YC(I2)))
IF(AD.GT.ABS(2.*R))AD=2.*ABS(R)
PANGLE=2.*ASIN(AD/(2.*R))
C      CALCULATE ANGLE BETWEEN 2 FARTHEST POINTS FROM
C      FARTHEST CORNER AND ADD TO PREVIOUS ANGLE.
AD=SQRT((XC(I3)-XC(I4))*(XC(I3)-XC(I4))
1  + (YC(I3)-YC(I4))*(YC(I3)-YC(I4)))
IF(AD.GT.ABS(2.*R))AD=2.*ABS(R)
PANGLE=2.*ASIN(AD/(2.*R))+PANGLE
C      ANGLE FOR THIS RADIUS IS NOW KNOWN, CALCULATE RISK
460 CONTINUE
ANAREA=PANGLE*R*STSIZE
APPROX=APPROX+ANAREA
NTOT=NRS(INGS)+1

```

```

      RATEI=RATE(INGS)*ANAREA*AREA(INGS,INSS)/AREA(INGS,NTOT)
C      CALCULATE CONTRIBUTION TO RISK
      DO 480 JJ=1,NLEI
      CALL RISK1(TI(JJ),R,INGS,RISK)
      IF (RISK-0.0000000001) 500,490,490
490   RSK(JJ)=RSK(JJ)+RISK*RATEI
480   CONTINUE
500   CONTINUE
      ARERR=((APPROX-AREA(INGS,INSS))/AREA(INGS,INSS))*100.
510   IF(ABS(ARERR).LE.ERRBND) GO TO 540
      WRITE (NWR,903) ARERR,INGS,INSS,SXNOT,SYNOT,NSTEPI
903   FORMAT(10X,"CAUTION: NUMERICAL INTEGRATION ERROR IN AREA IS ",
1     F8.2," % FOR (INSIDE) SOURCE ",2I3," LAT",F6.2," LONG",F8.2,
1     " NSTEPI=",I3)
      NSTEPI=NSTEPI*2
      IF(NSTEPI.GT.NSTEP MX) GO TO 10
      GO TO 3
10   LERR=.TRUE.
      RISKER=RISKER+COEF(INGS)*RSK(NLEI-2)/APPROX
540   DO 550 JJ=1,NLEI
      RSK(JJ)=COEF(INGS)*RSK(JJ)/APPROX
550   RSKTI(JJ)=RSKTI(JJ)+RSK(JJ)
      IF (JPRNT) 850,850,610
C      PRINT RISKS FOR THIS SOURCE.
610   WRITE(NWR,902) INGS,INSS,(RSK(I),I=1,NLEI)
902   FORMAT(" SOURCE",2I3," E(NO/YR): ",12E9.3)
      GO TO 850
C      ERROR PRINTOUT
800   WRITE (NWR,901) NERROR,INGS,INSS,IFAR,NPT,XNOT,YNOT,(XL(I),YL(I),
1     I=1,4),RC,RF,R,PANGLE,(XC(I),YC(I),I=1,4)
901   FORMAT (" ***** ERROR",I4," IN SUBROUTINE INSIDE. SOURCE NO.",
1     2I3," DEBUG VALUES FOLLOW.....",/10X,2I10,10(/10X,2F12.6))
      WRITE(NWR,1)SXNOT,SYNOT,NSTEPI
1     FORMAT(" LAT",F6.2," LONG",F8.2," NSTEPI=",I3)
      NSTEPI=NSTEPI*2
      IF(NSTEPI.LE.NSTEP MX)GO TO 3
      LERR2=.TRUE.
850   NSTEPI=NSTEPIS
      RETURN
      END
      SUBROUTINE NDTR(X,P,D)
C      X IS NO. OF STANDARDIZED NORMAL DEVIATES.
C      P IS COMP. CUMULATIVE VALUE (OUTPUT).
C      D IS DENSITY VALUE (OUTPUT)
      IF (X) 1,2,2
1     AX=-X
      GO TO 3
2     AX=X
3     IF (AX-6.0) 5,4,4
4     P=1.
      D=0.
      GO TO 6
5     T=1./(1.0+.2316419*AX)
      D=0.3989423*EXP(-X*X/2.0)
      P = 1.0 - D*T*((((1.330274*T - 1.821256)*T + 1.781478)*T -

```



```

1  0.3565638)*T + 0.3193815)
6  IF (X) 8,7,7
7  P=1.0-P
8  RETURN
END
SUBROUTINE NWNERK
C  INTERACTIVE VERSION FOR RISK CALCULATIONS IN THE NWNE REGION
C  DISC FILE INPUT OF ZONE DATA ON TAPE2
C  FOR A GIVEN LOCATION, WITH ROUTINE COMBINI, TO ADD RISKS FROM
C  SOURCES USING WESTERN ATTENUATIONS WITH RISKS FROM SOURCES USING
C  EASTERN ATTENUATIONS
C  DH WEICHERT MODIFIED FOR LAT. LONG. INPUT AND CONVERSION TO EASTING
C  AND NORTHINGS(Y) IN KM. THIS USE OF X + Y AGREES WITH MCGUIRES'S
C  MODIFIED BY FMA TO RUN VELOCITY AND ACCERATION DATA TOGETHER
C  R K MCGUIRE U.S.G.S. JANUARY 1975
C  PLANAR VERSION (CARTESIAN COORDINATES)
COMMON NRD,NWR,RSKTI(12)
COMMON C1,C2,C3,SIG,RZERO,RONE,AAA,BBB
COMMON NGS,NRS(23),AMO(24),AM1(24),LORS(24)
COMMON BETA(24),RATE(24),COEF(24),FDEPTH(24)
COMMON NSTEPO,NSTEP1
COMMON INDIC(4),AREA(23,11),X(23,11,2),Y(23,11,2)
COMMON/DEBG/SXNOT,SYNOT,ERRBND,NSTEPMX,LERR,RZ2,LERR2,RISKER
COMMON /MDATA/NSTEP,JPRNT,JPRNT2,JPRNT3,NLEI,II(12),RISKS(9),
1ATTEN(8,2,2)
COMMON/CNM/RNAME(6),SNAME(6),TIFS(8,2),RZ2S(23),JEW,NRG,SLAT,SLONG
1,IN,JN,NRGL
COMMON/LERRS/INCLUD,INCLD2,LNLEI,LRISKS,RKRATO
COMMON/SRSKC/SRSK(12,4)
DIMENSION BRISK(12)
LOGICAL LERR,INCLUD,LERR2,INCLD2,LNLEI,LRISKS
REWIND NRD$NRGS=NRG-1$DO 30 I=1,NRGS
31 READ(NRD)$IF(EOF(NRD))30,31
30 CONTINUE
DO 14 JW =1,2
IF(JW.EQ.1)GO TO 1
2 READ(NRD)
IF(EOF(NRD))1,2
1 READ(NRD) NGS,(NRS(I),I=1,NGS)
NGS1=NGS+1
DO 110 I=1,NGS1
READ(NRD) RZ2S(I),LORS(I),COEF(I),AMO(I),AM1(I),BETA(I),RATE(I)
1,FDEPTH(I),NAME
110 CONTINUE
DO 200 II=1,NGS
NRSII=NRS(II)+1
DO 150 JJ=1,NRSII
READ(NRD) X(II,JJ,1),Y(II,JJ,1),X(II,JJ,2),Y(II,JJ,2)
1,NAMES
CALL PRJC(X(II,JJ,1),Y(II,JJ,1))
CALL PRJC(X(II,JJ,2),Y(II,JJ,2))
C  DH WEICHERT CHANGES TO EASTING(X) + NORTHING(Y) IN KM. USING LAMBE
C  CONFORMAL PROJECTION
150 CONTINUE
200 CONTINUE

```

```

C      CALCULATE AREA OF EACH SUBSOURCE AND GROSS SOURCE.
      DO 400 II=1,NGS
      NTOT=NRS(II)+1
      AREA(II,NTOT)=0.0
      NAZIZ = NRS(II)
      DO 300 JJ=1,NAZIZ
      CALL AREAS(X(II,JJ,1),Y(II,JJ,1),X(II,JJ,2),Y(II,JJ,2),
1 X(II,JJ+1,1),Y(II,JJ+1,1),X(II,JJ+1,2),Y(II,JJ+1,2),AREA(II,JJ))
300 AREA(II,NTOT)=AREA(II,NTOT)+AREA(II,JJ)
      400 CONTINUE
      DO 424 J=1,2
      DO 90 I=1,NLEI
      90 BRISK(I)=0.0
      NSTEPI=NSTEP
      C1=ATTEN(1,J,JW)
      C2=ATTEN(2,J,JW)
      C3=ATTEN(3,J,JW)
      SIG=ATTEN(4,J,JW)
      RZERO=ATTEN(5,J,JW)
      RONE=ATTEN(6,J,JW)
      AAA=ATTEN(7,J,JW)
      BBB=ATTEN(8,J,JW)
      JAV=J+2*(JW-1)
      IF (BBB+0.00001) 101,102,102
      101 WRITE (JN ,924)
      924 FORMAT(// " INPUT ERROR: THE VALUE OF BBB MUST BE POSITIVE,"/
1 " BETWEEN 0.0 AND THE VALUE OF C2. EXECUTION STOPPED.")
      RETURN
      102 IF (C2-BBB) 101,103,103
C      IF BBB=0.0, SET EQUAL TO A SMALL NUMBER
      103 IF (BBB-0.00001) 104,105,105
      104 BBB=0.0000000001
C      COMPUTE BACKGROUND SEISMICITY
      105 IF(RATE(NGS1)-0.0000000001) 420,420,405
C      RBACK IS RADIUS OUT TO WHICH RISK FROM
C      BACKGROUND SEISMICITY IS CALCULATED.
      405 RBACK=150.
C      FOR BACKGROUND SEISMICITY, NSTEPI IS DOUBLED (AND
C      THEN HALVED AFTER CALCULATIONS).
      NSTEPI=2*NSTEPI
      CALL CIRCLE(RBACK,NGS1,1.,BRISK)
      NSTEPI=NSTEPI/2
      DO 410 I=1,NLEI
      410 BRISK(I)=COEF(NGS1)*BRISK(I)/10000.
      420 XNOT=SLAT$YNOT=-SLONG
      SXNOT=XNOT
      SYNOT=YNOT
      RISKER=0.
      LERR=.FALSE.
      LERR2=.FALSE.
      DO 450 I=1,NLEI
      450 RSKTI(I)=BRISK(I)
      CALL PRJC(XNOT,YNOT)
      DO 600 II=1,NGS
      NAZIZ = NRS(II)

```



```

      RZ2=RZ2S(II)
      DO 600 JJ=1,NAZIZ
      INGS=II
      INSS=JJ
      CALL RRISK(XNOT,YNOT,INGS,INSS,X(II,JJ,1),Y(II,JJ,1),X(II,JJ,
X2),Y(II,JJ,2),X(II,JJ+1,1),Y(II,JJ+1,1),X(II,JJ+1,2),Y(II,JJ+1,2))
600  CONTINUE
      IF(.NOT.LERR2.OR.INCLD2) GO TO 11
      WRITE(JN ,10)
10   FORMAT(/" ERROR IN SUBROUTINE INSIDE OR OUTSID"/)
      RETURN
11   IF(.NOT.LERR.OR.INCLUD) GO TO 12
      GDRISK=RSKTI(NLEI-2)-RISKER
      IF(GDRISK.LE.0.) GO TO 13
      RISKRR=RISKER/GDRISK
      IF(RISKRR.LT.RKRATO) GO TO 12
13   WRITE(JN ,15)      ERBND,RISKRR,RKRATO
15   FORMAT(/" ERROR IN AREA CALCULATION GREATER THAN ",F6.1," &"/
1" AND RATIO OF ERROR RISK CALC TO NON ERROR RISK CALC=",F6.2/
2" GREATER THAN ",F5.2/)
      RETURN
12   DO 610 I=1,NLEI
610  SRSK(I,JAV)=RSKTI(I)
424  CONTINUE
14   CONTINUE
      CALL COMBINI
      RETURN
      END
      SUBROUTINE OUTSID(XNOT,YNOT,INGS,INSS,XL,YL,XR,YR,AA,BB)
      COMMON NRD,NWR,RSKTI(12)
      COMMON C1,C2,C3,SIG,RZERO,RONE,AAA,BBB
      COMMON NGS,NRS(23),AMO(24),AM1(24),LORS(24)
      COMMON BETA(24),RATE(24),COEF(24),FDEPTH(24)
      COMMON NSTEPO,NSTEP1
      COMMON INDIC(4),AREA(23,11)
      COMMON/DEBG/SXNOT,SYNOT,ERRBND,NSTEPMX,LERR,RZ2,LERR2,RISKER
      COMMON /MATA/NSTEP,JPRNT,JPRNT2,JPRNT3,NLEI,TI(12),RISKS(9),
1ATTEN(8,2,2)
      DIMENSION XL(4),YL(4),XR(4),YR(4),AA(4),BB(4),XC(4),YC(4)
      DIMENSION RSK(12)
      LOGICAL LERR,LERR2
      NSTEPOS=NSTEP0
      SUBROUTINE FOR CALCULATING RISK WHEN SITE IS OUTSIDE
      (QUADRILATERAL) SOURCE AREA.
      DEFINE DISTANCE VALUES TO SELECT STEP SIZE.
      (RC IS CLOSEST DISTANCE BETWEEN SITE AND SOURCE.)
      RC BETWEEN 0.0 AND RZ1 IMPLIES STEP SIZE = NSTEPO.
      RC BETWEEN RZ1 AND RZ2 IMPLIES STEP SIZE = NSTEPO/2.
      RC BETWEEN RZ2 AND RZ3 IMPLIES LUMP RISK
      AT CENTER OF SOURCE (DEFINED BY AVERAGING LOCATIONS
      OF CORNER POINTS).
      RC GREATER THAN RZ3 IMPLIES IGNORE SOURCE.
      RZ1=100.
      RZ3=3000.
      TO BY-PASS THIS ALGORITHM, SET RZ1 TO A LARGE NUMBER.

```



```

C          TO PRODUCE 0-1 ALGORITHM (DISREGARD, OR CALCULATE USING
C          NSTEPO), SET RZ1=RZ2=RZ3=DISTANCE WITHIN WHICH YOU
C          WISH TO CONSIDER RISK.
C          FIND CLOSEST (IC,RC) AND FARTHEST (IFAR,RF) POINTS AND DISTANCES.
RC2=10000000000.
RF2=0.
DO 108 II=1,4
DIST=(XNOT-XL(II))*(XNOT-XL(II))+(YNOT-YL(II))*(YNOT-YL(II))
IF (RC2-DIST) 104,104,102
102 RC=SQRT(DIST)
RC2=DIST
IC=II
104 IF (DIST-RF2) 108,108,106
106 RF=SQRT(DIST)
RF2=DIST
IFAR=II
108 CONTINUE
ICS=0
C          SEE IF ANY SIDE LIES CLOSER THAN CLOSEST POINT
DO 150 II=1,4
C          IS SLOPE INFINITE?
IF(INDIC(II).EQ.1) GO TO 130
XS=XL(II)
YS=YNOT
GO TO 145
C          IS SLOPE ZERO?
130 IF (AA(II)-0.001) 131,140,140
131 IF (AA(II)+0.001) 140,140,132
132 XS=XNOT
YS=YL(II)
GO TO 145
C          SLOPE IS NOT ZERO, SO CALCULATE NEAREST POINT.
140 XS=(YNOT+(XNOT/AA(II))-BB(II))/(AA(II)+(1./AA(II)))
YS=((XNOT-XS)/AA(II))+YNOT
145 CALL BETWEN(XL(II),YL(II),XR(II),YR(II),XS,YS,INDIC(II),IANS)
IF (IANS) 150,146,148
146 NERROR=1
GO TO 800
148 DIST=(XNOT-XS)*(XNOT-XS)+(YNOT-YS)*(YNOT-YS)
IF (DIST-RC2) 149,150,150
149 RC2=DIST
RC=SQRT(DIST)
ICS=II
150 CONTINUE
3 APPROX=0.0
DO 290 II=1,NLEI
290 RSK(II)=0.
C          DETERMINE STEP SIZE FROM RZ1,RZ2, AND RZ3.
IF (RC-RZ1) 308,308,292
292 IF (RC-RZ2) 294,294,296
294 NSTEPX=NSTEPO/2
GO TO 310
296 IF (RC-RZ3) 298,298,850
C          IF RC IS BETWEEN RZ2 AND RZ3, CALCULATE RISK
C          ASSUMING SEISMICITY IS LUMPED AT CENTER (AVERAGE

```

```

C          OF CORNER POINTS).
298  XAVE=(XL(1)+XL(2)+XL(3)+XL(4))/4.
     YAVE=(YL(1)+YL(2)+YL(3)+YL(4))/4.
     R=SQRT((XAVE-XNOT)*(XAVE-XNOT)+(YAVE-YNOT)*(YAVE-YNOT))
     NTOT=NRS(INGS)+1
     RATEI=RATE(INGS)*AREA(INGS,INSS)/AREA(INGS,NTOT)
     DO 306 JJ=1,NLEI
     CALL RISK1(TI(JJ),R,INGS,RISK)
     IF (RISK-0.0000000001) 600,600,305
305  RSK(JJ)=COEF(INGS)*RISK*RATEI
306  RSKTI(JJ)=RSKTI(JJ) + RSK(JJ)
     GO TO 600
308  NSTEPX=NSTEPD
310  AN=NSTEPX
     STSIZE=(RF-RC)/AN
C          STEP THRU SOURCE AREA.
     DO 500 ISTEP=1,NSTEPX
     AI=ISTEP
     R=RC+(AI-0.5)*STSIZE
     NPT=0
     ANGLE=0.
     SIGNAL=1.
C          LOOP ON EACH SIDE
     DO 400 II=1,4
     IF(INDIC(II).EQ.1) GO TO 330
C          SIDE II IS A VERTICAL LINE
C          DOES CIRCLE (RADIUS R) INTERSECT IT?
     A=XL(II)-XNOT
     IF (A) 322,322,323
322  IF (R+A) 400,400,324
323  IF (R-A) 400,400,324
C          COMPUTE TWO INTERSECTION POINTS
324  X1=XL(II)
     B=SQRT(R*R-(X1-XNOT)*(X1-XNOT))
     Y1=YNOT+B
     X2=XL(II)
     Y2=YNOT-B
     GO TO 341
330  A=1.+AA(II)*AA(II)
     B=2.*(-XNOT+AA(II)*(BB(II)-YNOT))
     C=XNOT*XNOT+YNOT*YNOT+BB(II)*(BB(II)-2.*YNOT)-R*R
     D=B*B-4.*A*C
     IF (D) 400,400,340
C          TWO INTERSECTIONS, CALCULATE FIRST INTERSECTION POINT.
340  D=SQRT(D)
     X1=(-B+D)/(2.*A)
     Y1=AA(II)*X1 + BB(II)
     X2=(-B-D)/(2.*A)
     Y2=AA(II)*X2+BB(II)
C          SEE IF (X1,Y1) IS ON BOUNDARY
341  CALL BETWEN(XL(II),YL(II),XR(II),YR(II),X1,Y1,INDIC(II),IANS)
     IF (IANS) 360,342,345
342  NERROR=4
     GO TO 800
C          CALCULATE OTHER POINT, SEE IF IS ON BOUNDARY.

```

```

345 CALL BETWEN(XL(II),YL(II),XR(II),YR(II),X2,Y2,INDIC(II),IANS)
    IF (IANS) 348,346,350
346 NERROR=5
    GO TO 800
C    STORE FIRST POINT
348 NPT=NPT+1
    XC(NPT)=X1
    YC(NPT)=Y1
    GO TO 400
C    SEE IF THIS SIDE IS CLOSEST TO POINT, IF SO, TREAT SPECIALLY.
350 IF (II-ICS) 352,355,352
C    BOTH POINTS ARE ON BOUNDARY, CALCULATE ANGLE BETWEEN THEM.
352 SIGN=-1.
    GO TO 357
355 SIGN=1.
    SIGNAL=-1.
357 AD=SQRT((X1-X2)*(X1-X2)+(Y1-Y2)*(Y1-Y2))
    IF (AD-2.*R) 358,359,359
358 ANGLE=SIGN*2.*ASIN(AD/(2.*R)) + ANGLE
    GO TO 400
359 ANGLE=3.1415926536 +ANGLE
    GO TO 400
C    SEE IF SECOND POINT ONLY IS ON BOUNDARY
360 CALL BETWEN(XL(II),YL(II),XR(II),YR(II),X2,Y2,INDIC(II),IANS)
    IF (IANS) 400,362,370
362 NERROR=6
    GO TO 800
370 NPT=NPT+1
    XC(NPT)=X2
    YC(NPT)=Y2
400 CONTINUE
    IF(NPT.LT.1.OR.NPT.GT.4) GO TO 404
    GO TO (410,420,410,440),NPT
404 IF(SIGNAL)460,405,405
405 NERROR=7
    GO TO 800
410 NERROR=8
    GO TO 800
420 AD=SQRT((XC(1)-XC(2))*(XC(1)-XC(2))+(YC(1)-YC(2))*(YC(1)-YC(2)))
    IF(AD.GT.ABS(2.*R))AD=2.*ABS(R)
    ANGLE=ANGLE + SIGNAL*2.*ASIN(AD/(2.*R))
    GO TO 460
C    FOUR INTERSECTION POINTS (EACH ON A DIFFERENT SIDE).
C    DETERMINE ANGLE BY FINDING CLOSEST 2 INTERSECTIONS TO
C    FARTEST CORNER, CALCULATE ANGLE BETWEEN, AND ADD ANGLE
C    BETWEEN OTHER TWO INTERSECTIONS.
440 DIST1=10000000000.
    I1=0
    I2=0
    I3=0
    I4=0
    DO 450 JJ=1,4
    DIST=(XL(IFAR)-XC(JJ))*(XL(IFAR)-XC(JJ))
1  + (YL(IFAR)-YC(JJ))*(YL(IFAR)-YC(JJ))
    IF (DIST-DIST1) 442,444,444

```



```

442 DIST2=DIST1
    DIST1=DIST
    I4=I3
    I3=I2
    I2=I1
    I1=JJ
    GO TO 450
444 IF (DIST-DIST2) 445,446,446
445 DIST2=DIST
    I4=I3
    I3=I2
    I2=JJ
    GO TO 450
446 I4=I3
    I3=JJ
450 CONTINUE
C      CALCULATE ANGLE BETWEEN 2 CLOSEST POINTS TO FARTHEST CORNER.
    AD=SQRT((XC(I1)-XC(I2))*(XC(I1)-XC(I2))
1  + (YC(I1)-YC(I2))*(YC(I1)-YC(I2)))
    IF(AD.GT.ABS(2.*R))AD=2.*ABS(R)
    ANGLE=2.*ASIN(AD/(2.*R))
C      CALCULATE ANGLE BETWEEN 2 POINTS FARTHEST FROM
C      FARTHEST CORNER AND ADD TO PREVIOUS ANGLE.
    AD=SQRT((XC(I3)-XC(I4))*(XC(I3)-XC(I4))
1  + (YC(I3)-YC(I4))*(YC(I3)-YC(I4)))
    IF(AD.GT.ABS(2.*R))AD=2.*ABS(R)
    ANGLE=2.*ASIN(AD/(2.*R)) + ANGLE
460 CONTINUE
C      ANGLE FOR THIS RADIUS NOW KNOWN, CALCULATE RISK.
C      COMPUTE RATE OF EARTHQUAKES IN THIS ANNULAR SOURCE
    ANAREA=ANGLE*R*STSIZE
    APPROX=APPROX+ANAREA
    NTOT=NRS(INGS)+1
    RATEI=RATE(INGS)*ANAREA*AREA(INGS,INSS)/AREA(INGS,NTOT)
C      CALCULATE CONTRIBUTION TO RISK
    DO 480 JJ=1,NLEI
    CALL RISK1(TI(JJ),R,INGS,RISK)
    IF (RISK-0.0000000001) 500,490,490
490 RSK(JJ)=RSK(JJ)+RISK*RATEI
480 CONTINUE
500 CONTINUE
    ARERR=((APPROX-AREA(INGS,INSS))/AREA(INGS,INSS))*100.
510 IF(ABS(ARERR).LE.ERRBND) GO TO 540
    WRITE(NWR,903) ARERR,INGS,INSS,SXNOT,SYNOT,NSTEPD
903 FORMAT(10X,"CAUTION: NUMERICAL INTEGRATION ERROR IN AREA IS ",
1  F8.2," % FOR (OUTSIDE) SOURCE ",2I3," LAT",F6.2," LONG",F8.2,
1  " NSTEPO=",I3)
    NSTEPO=NSTEPD*2
    IF(NSTEPD.GT.NSTEPMX) GO TO 10
    GO TO 3
10 LERR=.TRUE.
    RISKE=RISKE+COEF(INGS)*RSK(NLEI-2)/APPROX
C      NORMALIZE BY COMPUTED (APPROXIMATE) AREA
540 DO 550 JJ=1,NLEI
    RSK(JJ)=COEF(INGS)*RSK(JJ)/APPROX

```

```

550 RSKTI(JJ)=RSKTI(JJ)+RSK(JJ)
600 IF (JPRNT) 850,850,610
C   PRINT RISKS FOR THIS SOURCE
610 WRITE(NWR,902) INGS,INSS,(RSK(I),I=1,NLEI)
902 FORMAT(" SOURCE",2I3," E(NO/YR): ",12E9.3)
    GO TO 850
C   ERROR PRINTOUT
800 WRITE(NWR,901)NERROR,INGS,INSS,IC,NPT,XNOT,YNOT,
    1(XL(I),YL(I),I=1,4),R,ANGLE,RC,RF,(XC(I),YC(I),I=1,4)
901 FORMAT(" ***** ERROR",I4," IN SUBROUTINE OUTSID. SOURCE NO.",2I3,
    1" DEBUG VALUES FOLLOW.....",/10X,2I5,5(/10X,4F14.6))
    WRITE(NWR,1)SXNOT,SYNOT,NSTEPO
    1 FORMAT(" LAT",F6.2," LONG",F8.2," NSTEPO=",I3)
    NSTEPO=NSTEPO*2
    IF(NSTEPO.LE.NSTEPMX.AND.NERROR.NE.1) GO TO 3
    LERR2=.TRUE.
850 NSTEPO=NSTEPOS
    RETURN
    END
    SUBROUTINE OUTFRM
    COMMON NRD,NWR,RSKTI(12)
    COMMON C1,C2,C3,SIG,RZERO,RONE,AAA,BBB
    COMMON NGS,NRS(23),AMO(24),AM1(24),LORS(24)
    COMMON BETA(24),RATE(24),COEF(24),FDEPTH(24)
    COMMON NSTEPO,NSTEPI
    COMMON INDIC(4),AREA(23,11),X(23,11,2),Y(23,11,2)
    COMMON /MDATA/NSTEP,JPRNT,JPRNT2,JPRNT3,NLEI,TI(12),RISKS(9),
    1ATTEN(8,2,2)
    COMMON/CNM/RNAME(6),SNAME(6),TIFS(8,2),RZ2S(23),JEW,NRG,SLAT,SLONG
    1,IN,JN,NRGL
    DIMENSION NCC(60)
    WRITE(NWR,1)
    1 FORMAT(1H1,T5,"ENERGY, MINES AND",T47,"ENERGIE, MINES ET"/
    1T5,"RESOURCES CANADA",T47,"RESOURCES CANADA"/
    2T5,"EARTH PHYSICS BRANCH",T47,"DIRECTION DE LA PHYSIQUE DU GLOBE"
    1//
    1///T5,"SEISMIC RISK CALCULATION *"
    4T47,"CALCULE DE RISQUE SEISMIQUE *"///)
    WRITE(NWR,2)
    2 FORMAT(//T5,"REQUESTED BY")
    DECODE(60,50,RNAME)NCC
50  FORMAT(60A1)
    DO 51 I=1,60$J=61-I$IF(NCC(J).EQ.10H                )GO TO 51$GO TO 52
51  CONTINUE
52  NBCH=(61-J)/2+20
    ENCODE(10,54,LINEF)NBCH,J
54  FORMAT(2H(T,I2,1H,,I2,3H1))
    WRITE(NWR,LINEF)(NCC(I),I=1,J)
    WRITE(NWR,12)
12  FORMAT(T5,"DEMANDE PAR"/)
    WRITE(NWR,3)
    3 FORMAT(//T5,"FOR SITE")
    DECODE(60,50,SNAME)NCC
    DO 55 I=1,60$J=61-I$IF(NCC(J).EQ.10H                )GO TO 55$GO TO 56
55  CONTINUE

```



```

56 NBCH=(61-J)/2+20
   ENCODE(10,54,LINEF)NBCH,J
   WRITE(NWR,LINEF)(NCC(I),I=1,J)
   WRITE(NWR,13)
13 FORMAT(T5,"POUR SITE"/)
   WRITE(NWR,4)SLAT,SLONG
   4 FORMAT(/T5,"LOCATED AT"/T29,F5.2," NORTH/NORD"
      2T49,F6.2," WEST/QUEST"/T5,"LOCATION"/)
   WRITE(NWR,5)(RISKS(J),J=1,4)
   5 FORMAT(/T5,"PROBABILITY OF EXCEEDENCE"/T5,"PER ANNUM"/
      235X,F4.2,6X,F5.3,5X,F8.6,6X,F5.3/
      3T5,"PROBABILITE DE"/T5,"DEPASSEMENT PAR ANNEE")
   WRITE(NWR,8)(TIFS(J,1),J=1,4)
   8 FORMAT(/ T5,"PEAK HORIZONTAL"/T5,"ACCELERATION (%G)"/
      135X,F4.1,7X,F4.1,8X,F4.1,8X,F4.1/
      2T5,"ACCELERATION HORIZONTAL"/T5,"MAXIMALE (%G)")
   WRITE(NWR,7)(TIFS(J,2),J=1,4)
   7 FORMAT(/ T5,"PEAK HORIZONTAL"/T5,"VELOCITY (CM/SEC)"/
      135X,F4.1,7X,F4.1,8X,F4.1,8X,F4.1/
      2T5,"VITESSE HORIZONTALE"/T5,"MAXIMALE (CM/SEC)")
   WRITE(NWR,10)
10 FORMAT(//////// T5,"* REFERENCE"/
      1T5,"NEW PROBABILISTIC STRONG SEISMIC GROUND"/
      2T5,"MOTION MAPS OF CANADA: A COMPILATION OF EARTHQUAKE"/
      3T5,"SOURCE ZONES, METHODS AND RESULTS"/
      4T5,"P.W. BASHAM, D.H. WEICHERT, F.M. ANGLIN, AND M.J. BERRY"/
      6T5,"EARTH PHYSICS BRANCH OPEN FILE NUMBER 82-   "/
      7T5,"OTTAWA, CANADA 1982")
   CALL DATE(IDATE)$CALL TIME(ITIME)$WRITE(NWR,20)IDATE,ITIME
20 FORMAT(60X,2A10)
   WRITE(NWR,9)
   9 FORMAT(////////)
   RETURN$END
   SUBROUTINE PRJC(RLAT1,RLONG1)
   COMMON/CTAS/SCALEK,CONE,TLONG,RO,RADCON
   DATA RAD,E,E2,A,ED2/0.01745329252,12.15482511000,0.00676865800218,
C637820640.1,0.041135927122/
   IF(RLAT1.GE.90.) GO TO 2
   P=RLAT1*RAD
   SP=SIN(P)
   RI=SCALEK*(COS(P)/(1.+SP)*((E+SP)/(E-SP))**ED2)**CONE
   QI=(TLONG-RLONG1)*RADCON
   RLAT1=-RI*SIN(QI)
   RLONG1=RO-RI*COS(QI)
C   NOTE THE TRANSPOSED INTERPRETATION OF THE ARGUMENT SEQUENCE.
C   THE FIRST ARG, LAT. IS NOW X, SECOND Y
   RETURN
   2 RLAT1=1.E8
   RETURN
   END
   SUBROUTINE RISK1(TIC,REPIS,INGS,RISK)
C   D WEICHERT S VERSION MAY 1981
C   MODIFIED MAY 28
   COMMON NRD,NWR,RSKTI(12)
   COMMON C1,C2,C3,SIG,RZERO,RONE,AAA,BBB

```



```

70 IF (AMZ-AMSTAR) 80,75,75
C   ALL OF MAGNITUDE INTEGRATION LIES ABOVE AMSTAR.
75 CALL ERISK(AMZ,AM1J      ,AAA,BBB,O.,RLN,SIGG,BETAJ      ,TIC,
1   G1,G2,G3,G4,CON1,CON2,CON3)
77 RISK=((1.-AK)*G1 + AK*G2 + AK*(G3-G4)*CON1*CON2+AK*SPIKE*PHISTR)*
1ANEQ
GO TO 100
C   SOME OF MAGNITUDE INTEGRATION LIES ABOVE MSTAR, SOME BELOW.
80 CALL ERISK(AMZ,AMSTAR,C1,C2,C3,RLN,SIGG,BETAJ      ,TIC,
1   G1,G2,G3,G4,CON1,CON2,CON3)
CALL ERISK(AMSTAR,AM1J      ,AAA,BBB,O.,RLN,SIGG,BETAJ      ,TIC,
1   GG1,GG2,GG3,GG4,CCON1,CCON2,CCON3)
RISK=((1.-AK)*G1 +AK*G2 +AK*(G3-G4)*CON1*CON2
1 + (1.-AK)*(GG1-GG2) + AK*(GG3-GG4)*CCON1*CCON2
2*EXP(BETAJ*(AMZ-AMSTAR))+AK*SPIKE*PHISTR)*ANEQ
100 RETURN
END
SUBROUTINE RRISK(XNOT,YNOT,INGS,INSS,
1 X1,Y1,X2,Y2,X3,Y3,X4,Y4)
COMMON NRD,NWR,RSKTI(12)
COMMON C1,C2,C3,SIG,RZERO,RONE,AAA,BBB
COMMON NGS,NRS(23),AMO(24),AM1(24),LORS(24)
COMMON BETA(24),RATE(24),COEF(24),FDEPTH(24)
COMMON NSTEPO,NSTEPI
COMMON INDIC(4),AREA(23,11)
COMMON /MDATA/NSTEP,JPRNT,JPRNT2,JPRNT3,NLEI,TI(12),RISKS(9),
1ATTEN(8,2,2)
DIMENSION XL(4),YL(4),XR(4),YR(4),AA(4),BB(4)
C   SUBROUTINE WHICH LOADS TEMPORARY ARRAYS WITH THIS
C   SUBSOURCE'S CORNERS AND DETERMINES IF THIS SITE IS
C   WITHIN OR WITHOUT THE SUBSOURCE.
XL(1)=X1
YL(1)=Y1
XR(1)=X2
YR(1)=Y2
XL(2)=X2
YL(2)=Y2
XR(2)=X4
YR(2)=Y4
XL(3)=X3
YL(3)=Y3
XR(3)=X1
YR(3)=Y1
XL(4)=X4
YL(4)=Y4
XR(4)=X3
YR(4)=Y3
C   DETERMINE IF ANY SIDES ARE VERTICAL LINES
DO 200 II=1,4
DIF=XL(II)-XR(II)
IF (DIF) 140,180,160
140 IF (DIF+0.01) 190,190,180
160 IF (DIF-0.01) 180,190,190
C   INDIC(II)=1 IMPLIES NOT A VERTICAL LINE
C   INDIC(II)=2 IMPLIES A VERTICAL LINE (INFINITE SLOPE).

```



```

180  INDIC(II)=2
    AA(II)=0.
    BB(II)=0.
    GO TO 200
190  INDIC(II)=1
    AA(II)=(YL(II)-YR(II))/(XL(II)-XR(II))
    BB(II)=YL(II)-AA(II)*XL(II)
C      AA(II) IS SLOPE OF II'TH SIDE
C      BB(II) IS INTERCEPT OF II'TH SIDE
200  CONTINUE
C      DETERMINE IF SITE IS INSIDE SOURCE AREA.
    DO 220 II=1,4
    IJ=5-II
    IF(INDIC(II).GE.2) GO TO 215
    DIFNOT=YNOT-AA(II)*XNOT-BB(II)
    DIF=YL(IJ)-AA(II)*XL(IJ)-BB(II)
211  IF (DIF) 214,214,212
212  IF (DIFNOT) 400,400,220
214  IF (DIFNOT) 220,400,400
215  DIF=XL(IJ)-XL(II)
    DIFNOT=XNOT-XL(II)
    GO TO 211
220  CONTINUE
C      IF DO LOOP FINISHED, POINT LIES WITHIN AREA.
    CALL INSIDE(XNOT,YNOT,INGS,INSS,XL,YL,XR,YR,AA,BB)
    GO TO 900
400  CALL OUTSID(XNOT,YNOT,INGS,INSS,XL,YL,XR,YR,AA,BB)
900  RETURN
    END
    SUBROUTINE SHOUT
    COMMON /MDATA/NSTEP,JPRNT,JPRNT2,JPRNT3,NLEI,TI(12),RISKS(9),
1ATTEN(8,2,2)
    COMMON/CNM/RNAME(6),SNAME(6),TIFS(8,2),RZ2S(23),JEW,NRG,A,B
1,IN,JN,NRGL
    WRITE(JN,4)A,B
4  FORMAT(// " SITE ",F7.3,"N",F9.3,"W"/)
    WRITE(JN,1)(RISKS(J),J=1,4)
1  FORMAT(" PROB",4F10.6/)
    WRITE(JN,2)(TIFS(J,1),J=1,4)
2  FORMAT(" ACCL",4F10.2/)
    WRITE(JN,3)(TIFS(J,2),J=1,4)
3  FORMAT(" VELC",4F10.2//)
    RETURN$END
    SUBROUTINE SRISK
C      INTERACTIVE VERSION FOR RISK CALCULATIONS AT SPECIFIC SITES
C      DISC FILE INPUT OF ZONE DATA ON TAPE2
C      PRINTED OUTPUT ON TAPE6
C      MAINROUTINE
C      DH WEICHERT MODIFIED FOR LAT. LONG. INPUT AND CONVERSION TO EASTIN
C      AND NORTHINGS(Y) IN KM. THIS USE OF X + Y AGREES WITH MCGUIRES"S
C      MODIFIED BY FMA TO RUN VELOCITY AND ACCERATION DATA TOGETHER
C      R K MCGUIRE U.S.G.S. JANUARY 1975
C      PLANAR VERSION (CARTESIAN COORDINATES)
C
    COMMON NRD,NWR,RSKTI(12)

```



```

COMMON C1,C2,C3,SIG,RZERO,RONE,AAA,BBB
COMMON NGS,NRS(23),AMO(24),AM1(24),LORS(24)
COMMON BETA(24),RATE(24),COEF(24),FDEPTH(24)
COMMON NSTEPO,NSTEP1
COMMON INDIC(4),AREA(23,11),X(23,11,2),Y(23,11,2)
COMMON/DEBG/SXNOT,SYNOT,ERRBND,NSTEPMX,LERR,RZ2,LERR2,RISKER
COMMON /MDATA/NSTEP,JPRNT,JPRNT2,JPRNT3,NLEI,TI(12),RISKS(9),
1ATTEN(8,2,2)
COMMON/CNM/RNAME(6),SNAME(6),TIFS(8,2),RZ2S(23),JEW,NRG,SLAT,SLONG
1,IN,JN,NRGL
COMMON/LERRS/INCLUD,INCLD2,LNLEI,LRISKS,RKRATO
DIMENSION BRISK(12),TIF(8)
LOGICAL LERR,INCLUD,LERR2,INCLD2,LNLEI,LRISKS
IF(NRG.EQ.NRGL)GO TO 1$NRGL=NRG
REWIND NRD
C SKIP TO APPROPRIATE FILE
IF(NRG.LE.1) GO TO 712$NRGS=NRG-1$DO 710 I=1,NRGS
711 READ(NRD)$IF(EOF(NRD))710,711
710 CONTINUE
712 READ(NRD) NGS,(NRS(I),I=1,NGS)
NGS1=NGS+1
DO 110 I=1,NGS1
110 READ(NRD) RZ2S(I),LORS(I),COEF(I),AMO(I),AM1(I),BETA(I),RATE(I)
1,FDEPTH(I),NAME
DO 200 II=1,NGS
NRSII=NRS(II)+1
DO 150 JJ=1,NRSII
READ(NRD) X(II,JJ,1),Y(II,JJ,1),X(II,JJ,2),Y(II,JJ,2)
1,NAMES
CALL PRJC(X(II,JJ,1),Y(II,JJ,1))
CALL PRJC(X(II,JJ,2),Y(II,JJ,2))
C OH WEICHERT CHANGES TO EASTING(X) + NORTHING(Y) IN KM. USING LAMBE
C CONFORMAL PROJECTION
150 CONTINUE
200 CONTINUE
C CALCULATE AREA OF EACH SUBSOURCE AND GROSS SOURCE.
DO 400 II=1,NGS
NTOT=NRS(II)+1
AREA(II,NTOT)=0.0
NAZIZ = NRS(II)
DO 300 JJ=1,NAZIZ
CALL AREAS(X(II,JJ,1),Y(II,JJ,1),X(II,JJ,2),Y(II,JJ,2),
1 X(II,JJ+1,1),Y(II,JJ+1,1),X(II,JJ+1,2),Y(II,JJ+1,2),AREA(II,JJ))
300 AREA(II,NTOT)=AREA(II,NTOT)+AREA(II,JJ)
400 CONTINUE
1 DO 424 J=1,2
NSTEPO=NSTEP1=NSTEP
DO 90 I=1,NLEI
90 BRISK(I)=0.0
C1=ATTEN(1,J,JEW)
C2=ATTEN(2,J,JEW)
C3=ATTEN(3,J,JEW)
SIG=ATTEN(4,J,JEW)
RZERO=ATTEN(5,J,JEW)
RONE=ATTEN(6,J,JEW)

```

```

      AAA=ATTEN(7,J,JEW)
      BBB=ATTEN(8,J,JEW)
      IF (BBB+0.00001) 101,102,102
101  WRITE (JN ,924)
924  FORMAT(// " INPUT ERROR:  THE VALUE OF BBB MUST BE POSITIVE,"/
1    " BETWEEN 0.0 AND THE VALUE OF C2. EXECUTION STOPPED.")
      RETURN
102  IF (C2-BBB) 101,103,103
C    IF BBB=0.0, SET EQUAL TO A SMALL NUMBER
103  IF (BBB-0.00001) 104,105,105
104  BBB=0.0000000001
105  CONTINUE
C    COMPUTE BACKGROUND SEISMICITY
      IF(RATE(NGS1)-0.0000000001) 420,420,405
C    RBACK IS RADIUS OUT TO WHICH RISK FROM
C    BACKGROUND SEISMICITY IS CALCULATED.
405  RBACK=150.
C    FOR BACKGROUND SEISMICITY, NSTEPI IS DOUBLED (AND
C    THEN HALVED AFTER CALCULATIONS).
      NSTEPI=2*NSTEPI
      CALL CIRCLE(RBACK,NGS1,1.,BRISK)
      NSTEPI=NSTEPI/2
      DO 410 I=1,NLEI
410  BRISK(I)=COEF(NGS1)*BRISK(I)/10000.
420  YNOT=-SLONG$XNOT=SLAT
      SXNOT=XNOT
      SYNOT=YNOT
      RISKER=0.
      LERR=.FALSE.
      LERR2=.FALSE.
      DO 450 I=1,NLEI
450  RSKTI(I)=BRISK(I)
      CALL PRJC(XNOT,YNOT)
      DO 600 II=1,NGS
      NAZIZ = NRS(II)
      RZ2=RZ2S(II)
      DO 500 JJ=1,NAZIZ
      INGS=II
      INSS=JJ
      CALL RRISK(XNOT,YNOT,INGS,INSS,X(II,JJ,1),Y(II,JJ,1),X(II,JJ,
X2),Y(II,JJ,2),X(II,JJ+1,1),Y(II,JJ+1,1),X(II,JJ+1,2),Y(II,JJ+1,2))
500  CONTINUE
600  CONTINUE
      IF(.NOT.LERR2.OR.INCLD2) GO TO 11
      WRITE(JN ,10)
10  FORMAT(// " ERROR IN SUBROUTINE INSIDE OR OUTSID")
      RETURN
11  IF(.NOT.LERR.OR.INCLUD) GO TO 12
      GDRISK=RSKTI(NLEI-2)-RISKER
      IF(GDRISK.LE.0.) GO TO 13
      RISKRR=RISKER/GDRISK
      IF(RISKRR.LT.RKRATO) GO TO 12
13  WRITE(JN ,15)      ERRBND,RISKRR,RKRATO
15  FORMAT(// " ERROR IN AREA CALCULATION GREATER THAN ",F6.1," %"/
1    " AND RATIO OF ERROR RISK CALC TO NON ERROR RISK CALC=",F6.2/

```

```

      2" GREATER THAN ",F5.2/)
      RETURN
12  CONTINUE
      DO 620 I=1,NLEI
620  RSKTI(I)=1.-EXP(-RSKTI(I))
C    ESTIMATE INTENSITIES AT RISKS DESIRED.
      RISKS(9)=0.0
      IA=0
      IF(RISKS(1)-0.0000000001) 700,700,625
625  DO 630 IRK=1,8
      IF (RISKS(IRK)-RSKTI(1)) 640,640,630
630  TIF(IRK)=1000000.
      GO TO 700
640  IA=IA+1
      IF (IA-NLEI) 650,645,645
645  TIF(IRK)=1000000.
      IRK=IRK+1
      IF (RISKS(IRK)-0.0000000001)680,680,645
650  IF(RISKS(IRK)-RSKTI(IA+1))640,655,655
655  CONTINUE
      TIF(IRK)=(ALOG(RSKTI(IA)/RISKS(IRK)))
      1 /((ALOG(RSKTI(IA)/RSKTI(IA+1)))
      TIF(IRK)=TI(IA)+TIF(IRK)*(TI(IA+1)-TI(IA))
      IRK=IRK+1
      IF (RISKS(IRK)-0.0000000001)680,680,660
660  IF(RISKS(IRK)-RSKTI(IA+1)) 640,655,655
680  IRK=IRK-1
      DO 685 I=1,IRK
      IF (TIF(I)-999999.)683,685,685
683  TIF(I)=EXP(TIF(I))
685  CONTINUE
      700 DO 722 I=1,IRK
      722 TIFS(I,J)=TIF(I)
      424 CONTINUE
      RETURN$END

```


APPENDIX CRevised Parameters for Eastern Canadian
and some Northeastern U.S. Earthquakes

For the preparation of the earthquake source models of eastern Canada by Basham et al. (1979), a review was made of most of the pre-1968 earthquakes in the region that had previously catalogued magnitudes of 4 or greater. Revised parameters with magnitudes quoted to the nearest half-magnitude category were listed in their Table 1. It is the purpose of this appendix to document the reasons for the changes that were made, particularly in magnitude, so that corresponding changes can be made to the master Canadian Earthquake Epicentre File (CEEFF) maintained by the Earth Physics Branch.

The review of information available for most of the earthquakes was not exhaustive and in many, if not most, of the cases a further review would be justified to either confirm or adjust the revised parameters. However, the changes that have been made are considered a significant improvement on previously catalogued data for the purposes, in both Basham et al. (1979) and this report, of estimating magnitude recurrence relations for eastern Canadian earthquake source zones.

The two general categories of review were the following. During the first preparation of the CEEFF the magnitudes of most of the historical earthquakes were based on Smith's (1962, 1966) epicentral intensities. Comprehensive review of macroseismic information for a number of earthquakes has shown that many of Smith's epicentral intensities were based on exaggerated effects or were not representative of the epicentral region. For

the additional earthquakes treated here, this information has been reviewed and revisions made to magnitude if a change of at least one-half unit is appropriate. For many of the smaller earthquakes the original sources of information have not been reviewed; the revisions are based solely on Smith's summary description.

For many earthquakes in the time period 1935 to 1967, the original amplitude data, used to compute M_L which appears in the catalogues and on the CEEF, was available. These data have been used directly to compute $m_b(Lg)$ and magnitude has been revised if a change of at least one magnitude category is appropriate. The original seismograms were not used to check the original amplitude data.

Basham et al. (1979; their Table 1) listed earthquakes with magnitude categories (i.e., to the nearest half unit) of 4 or greater. Thus, earthquakes with revisions that resulted in magnitude categories less than 4 did not appear in that list. All such revisions are included in the following table. In those cases for which the revised magnitude is a recomputed $m_b(Lg)$, the magnitude is quoted to the nearest tenth of a unit (although it should not be considered this accurate) and the remark is a simple statement " $m_b(Lg)$ from n stations." It should be noted that many of the earthquakes in the following table do not appear in Appendix A, the individual source zone lists, because they do not pass the completeness test used for magnitude recurrence calculations.

In the following table the first entry for each earthquake gives the parameters currently on the CEEF, the second entry the revision. This is followed by a brief explanation for the change.

No.	Year	M	D	H	M	Lat.(°N)	Long.(°W)	Mag.
1.	1663	02	05	17	30	47.6	70.1	7.7 7.0

For this earthquake and three later earthquakes in the Charlevoix zone described below (Nos. 3, 10 and 13), revised magnitude estimates are made on the basis of comparison of intensity information with the well-defined intensity and isoseismal data available for the 1925, M7 earthquake. For this event the intensity effects in the distance range 100-1000 km (i.e., ignoring the near-in landslide phenomena that are not a good indication of earthquake size) are very similar to those of 1925, with only a slight tendency to larger intensities. Therefore the earthquake has been assigned M7.

2.	1665	02	24			47.8	70.0	6.4 5.5
----	------	----	----	--	--	------	------	------------

The Smith (1962) epicentral intensity, from which magnitude 6.4 was derived, is considered to be an over-estimate. Reports of low intensities in New England suggest a magnitude of 5.5 or smaller.

3.	1791	12	06	20		47.4	70.5	6.3 6.0
----	------	----	----	----	--	------	------	------------

A comparison of the well-defined intensities for this earthquake with those of 1925 suggests that it is approximately one magnitude unit smaller; it has therefore been assigned M6.

4.	1817	05	22	20		46.0	69.0	5.7 5.0
----	------	----	----	----	--	------	------	------------

The evidence for Smith's (1962) epicentral intensity is poor. The felt area, which translates to M5, is considered to provide a better estimate of magnitude.

5.	1831	07	14			47.6	70.1	5.7 5.0
----	------	----	----	--	--	------	------	------------

Very little intensity information is available. The magnitude is reduced to 5 on the assumption that Smith's (1962) epicentral intensity represents the results of poor construction and not general effects in the epicentral region.

6.	1840	09	10			43.2	79.9	4.4 4.0
----	------	----	----	--	--	------	------	------------

Described as "a violent shock" at Hamilton, this earthquake is most likely a shallow event of M4 or smaller, similar to such events in more recent years.

No.	Year	M	D	H	M	Lat.(°N)	Long.(°W)	Mag.
7.	1853	03	12	07		43.7	75.5	5.0 4.5

Felt area not extensive and more consistent with M4.5.

8.	1853	03	13	10		43.1	79.4	4.4 4.0
----	------	----	----	----	--	------	------	------------

Felt area more consistent with M4.

9.	1857	12	23			44.1	70.2	5.7 4.5
----	------	----	----	--	--	------	------	------------

No evidence for high epicentral intensity or extensive felt area.

10.	1860	10	17	11	15	47.5	70.1	6.7 6.0
-----	------	----	----	----	----	------	------	------------

A comparison of the well-defined intensities of this earthquake with those of 1925 suggests that it is approximately one magnitude unit smaller; it has therefore been assigned M6.

11.	1861	07	12			45.4	75.4	5.7 5.0
-----	------	----	----	--	--	------	------	------------

Felt area (Montreal, Ottawa, Odgensburg) suggests M5 or smaller; chimney damage at Ottawa not considered representative of high epicentral intensity.

12.	1869	12				47.5	70.5	4.4 4.0
-----	------	----	--	--	--	------	------	------------

Felt information appropriate to M4 or smaller.

13.	1870	10	20	16	30	47.4	70.5	7.0 6.5
-----	------	----	----	----	----	------	------	------------

A comparison of the well-defined intensities of this earthquake with those of 1925 suggests that it is approximately one-half magnitude unit smaller; it has therefore been assigned M6.5

14.	1871	01	09			47.5	70.1	4.4 4.0
-----	------	----	----	--	--	------	------	------------

Felt information consistent with M4 or smaller.

15.	1872	01	09			47.5	70.5	5.7 5.0
-----	------	----	----	--	--	------	------	------------

Felt area, suggesting M5, is considered more representative than Smith's (1962) epicentral intensity.

No.	Year	M	D	H	M	Lat.(°N)	Long.(°W)	Mag.
16.	1873	07	06	14	30	43.0	79.5	5.0 4.5

Felt area consistent with M4.5 or smaller.

17.	1874	02	27			44.8	68.7	4.4 4.0
-----	------	----	----	--	--	------	------	------------

Felt area consistent with M4 or smaller.

18.	1887	05	27	06	15	47.5	70.5	4.3 4.0
-----	------	----	----	----	----	------	------	------------

Felt area consistent with M4 or smaller.

19.	1896	03	22			45.2	67.2	4.4 4.0
-----	------	----	----	--	--	------	------	------------

Felt area consistent with M4.

20.	1897	03	23			45.5	73.6	5.7 5.0
-----	------	----	----	--	--	------	------	------------

No evidence for high epicentral intensity. Felt area suggests much smaller event; M5 adopted as a compromise.

21.	1897	05	27			44.5	73.5	5.0 4.5
-----	------	----	----	--	--	------	------	------------

Felt area quoted by Smith (1962) (150,000 mi²) is not consistent with "felt from Montreal to Burlington, Vt.", which suggests M4.5 or smaller.

22.	1906	06	27			41.4	81.6	4.4 4.0
-----	------	----	----	--	--	------	------	------------

No evidence for M larger than 4.

23.	1906	10	20			43.8	68.8	4.4 4.0
-----	------	----	----	--	--	------	------	------------

No evidence for M larger than 4.

24.	1908	05	14	04	45	44.0	65.8	4.3 4.0
-----	------	----	----	----	----	------	------	------------

Felt area consistent with M4.

25.	1909	12	19	20		46.5	60.5	5.0 4.0
-----	------	----	----	----	--	------	------	------------

There is little evidence to support Smith's (1962) epicentral intensity. The felt area, "throughout Cape Breton", is more consistent with M4.

No.	Year	M	D	H	M	Lat.(°N)	Long.(°W)	Mag.
26.	1910	01	23	01	30	43.8	70.4	3.7 4.0

Report of "articles thrown from shelves" suggests magnitude greater than 3.7

27.	1910	10	25	09	30	47.6	69.8	4.3 4.0
-----	------	----	----	----	----	------	------	------------

Felt area consistent with M4.

28.	1912	12	11	10	15	45.0	68.0	3.7 4.0
-----	------	----	----	----	----	------	------	------------

Felt area suggests magnitude 4.

29.	1914	02	10	18	31	45.0 46.0	76.9 75.0	5.5
-----	------	----	----	----	----	--------------	--------------	-----

Klotz (1915) determined that this earthquake was felt over an area of 500,000 km², which is compatible with the magnitude (M_L) calculated by Smith (1962) from Ottawa data. Klotz chose an epicentre northeast of Ottawa, his paper implying that the Ottawa seismograms unambiguously indicated a northeast direction. Smith determined an epicentre southwest of Ottawa using recorded arrival times at Ottawa, Harvard and Ithaca. He did not use the Toronto arrival time which, in combination with Ottawa and Harvard, supports an epicentre northeast of Ottawa. The Ithaca arrival times are apparently no longer available (A.E. Stevens, unpublished notes, 1976). The revision is an epicentre northeast of Ottawa, which places this earthquake in the region north of the Ottawa River that has been most active in recent years, rather than in an essentially aseismic region of eastern Ontario. The revised epicentre must be considered to have a large uncertainty.

30.	1914	02	22	19	15	45.0	70.5	4.4 4.0
-----	------	----	----	----	----	------	------	------------

Felt area consistent with M4.

31.	1915	07	27	16	30	44.0	65.0	4.3 4.0
-----	------	----	----	----	----	------	------	------------

Felt information consistent with M4 or smaller.

32.	1916	01	05	13	56	43.7	73.7	4.4 4.0
-----	------	----	----	----	----	------	------	------------

Felt area consistent with M4 or smaller.

No.	Year	M	D	H	M	Lat.(°N)	Long.(°W)	Mag.
33.	1916	04	24	16	07	47.0	77.0	4.3 4.0

Recorded by Ottawa seismograph but not instrumentally located and not reported felt. M reduced to 4 to give this earthquake less weight because of epicentral uncertainty.

34.	1917	06	12	02	00	49.0	68.0	4.3 4.0
-----	------	----	----	----	----	------	------	------------

Felt area more consistent with M4.

35.	1918	08	21	04	20	44.2	70.6	5.7 4.5
-----	------	----	----	----	----	------	------	------------

Felt area suggests M4.5 or lower. "Damaged chimneys" not sufficient for Smith's (1962) epicentral intensity.

36.	1925	03	07	02	30	47.8	69.8	4.4 4.0
-----	------	----	----	----	----	------	------	------------

Felt area consistent with M4.

37.	1925	10	09	14	00	43.7	71.1	4.3 4.0
-----	------	----	----	----	----	------	------	------------

Felt area consistent with M4.

38.	1926	08	28	21	30	44.7	70.0	4.4 4.0
-----	------	----	----	----	----	------	------	------------

Felt area consistent with M4.

39.	1927	07	25	00	56	47.3	71.0	4.3 4.0
-----	------	----	----	----	----	------	------	------------

Felt area consistent with M4.

40.	1928	02	08			45.3	69.0	5.0 4.5
-----	------	----	----	--	--	------	------	------------

No evidence to suggest M greater than 4.5.

41.	1928	04	25	23	38	44.5	71.2	4.3 4.0
-----	------	----	----	----	----	------	------	------------

No evidence to suggest magnitude greater than 4.

No.	Year	M	D	H	M	Lat.(°N)	Long.(°W)	Mag.
42.	1929	08	12	11	24	42.9	78.4	5.8 5.5

The original magnitude by Smith (1966) was M_L . The felt area suggests a magnitude of about 5. Street and Trucotte (1977) derived $m_b(Lg)$ 5.2. We have adopted $M_{5.5}$ and leave more definitive work to our U.S. colleagues.

43.	1931	04	20	19	54	43.4	73.7	5.0 4.5
-----	------	----	----	----	----	------	------	------------

$M_{L5.0}$ was determined from one station and is reduced by one-half magnitude unit assuming similarity with recomputations of $m_b(Lg)$ described below.

44.	1934	10	29	20	07	42.2	80.2	4.3 4.0
-----	------	----	----	----	----	------	------	------------

There is no evidence for Smith's (1966) high epicentral intensity (V).

45.	1935	11	01	17	02	46.8	79.1	4.6 4.1
-----	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 1 station.

46.	1935	11	02	00	42	46.8	79.1	4.7 4.2
-----	------	----	----	----	----	------	------	------------

$m_b(LG)$ from 1 station.

47.	1935	11	02	14	31	47.2	78.2	5.4 4.9
-----	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 1 station.

48.	1935	11	05	10	10	46.8	79.1	4.5 3.9
-----	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 1 station.

49.	1935	11	25	06	19	46.8	79.1	4.7 4.1
-----	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 1 station.

50.	1935	11	27	19	31	46.8	79.1	4.6 4.1
-----	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 1 station.

No.	Year	M	D	H	M	Lat.(°N)	Long.(°W)	Mag.
51.	1936	01	20	06	01	46.8	79.1	4.5 3.8
$m_b(Lg)$ from 1 station.								
52.	1936	03	25	01	27	46.8	79.1	4.6 4.0
$m_b(Lg)$ from 1 station.								
53.	1938	05	17	18	32	49.0	68.0	4.6 3.9
$m_b(Lg)$ from 1 station.								
54.	1939	06	24	17	20	47.3	70.4	4.8 4.5
Felt area consistent with M4.5. $m_b(Lg)$ from 1 station is 4.4.								
55.	1939	10	19	11	53	47.8	69.8	5.8 5.6
The original magnitude is M_L at 1 station. Felt area gives a lower limit of M5.5. Street and Turcotte (1977) calculated $m_b(Lg)$ 5.6, which is accepted.								
56.	1939	10	27	01	36	47.8	69.8	5.2 4.5
$m_b(Lg)$ from 1 station is 4.8. Intensity information is sparse. Basham et al. (1979) assigned M4.5 as more representative of an aftershock of the 19 October, M5.8 mainshock.								
57.	1939	11	07	02	40	47.8	70.5	4.3 4.1
$m_b(Lg)$ from 1 station.								
58.	1940	12	20	07	27	43.8	71.3	5.8 5.0
59.	1940	12	24	13	43	43.8	71.3	5.8 5.0

Felt areas for these two events are consistent with M5. More definitive reassessments that may be available from U.S. studies have not been searched.

No.	Year	M	D	H	M	Lat.(°N)	Long.(°W)	Mag.
60.	1942	08	26	17	54	46.8	77.5	4.1 3.7

$m_b(Lg)$ from 1 station.

61.	1942	09	11	11	05	49.2	67.4	4.4 3.7
-----	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 1 station.

62.	1943	01	14	21	32	45.3	69.6	5.4 5.0
-----	------	----	----	----	----	------	------	------------

M_L reduced by one-half magnitude unit. Intensity data have not been reassessed.

63.	1944	04	09	12	44	49.9	67.4	5.4 4.9
-----	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 2 stations.

64.	1944	09	05	04	38	45.0	74.9	5.9 5.6
-----	------	----	----	----	----	------	------	------------

The original magnitude was from Gutenberg and Richter (1954) class "d" (5.3 to 5.9) from which Smith (1966) selected the maximum in the range, 5.9, believing that data were available suggesting at least this magnitude. The number of stations reporting this earthquake in the International Seismological Summary suggests a magnitude less than 5.9. Examination of epicentral effects, with due consideration to design and construction techniques and to soil conditions suggests an epicentral intensity of VII, corresponding to magnitude 5.7. Consideration of seismograph data available from Canadian stations suggests m_b in the range 5.4 to 5.7 and M_L 5.6 ± 0.3 ; the latter is adopted here. (From A.E. Stevens, unpublished notes, 1976; more details available on request.) Street and Turcotte (1977) subsequently determined $m_b(Lg)$ 5.8.

65.	1944	11	05	19	07	48.7	80.8	5.1 4.4
-----	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 3 stations.

66.	1945	10	09	13	18	47.8	69.8	4.9 4.7
-----	------	----	----	----	----	------	------	------------

Smith's (1966) M_L magnitude was determined from 3 stations, one of which (Ottawa) was at large distance. An average M_L from the nearest two stations is 4.7.

No.	Year	M	D	H	M	Lat.(°N)	Long.(°W)	Mag.
77.	1952	01	30	04	00	44.5	73.2	5.0 4.5

Felt only locally, suggesting M4.5 or smaller.

78.	1952	03	30	13	11	47.6	69.9	4.4 4.1
-----	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 2 stations.

79.	1952	08	25	00	07	43.0	74.5	4.3 4.0
-----	------	----	----	----	----	------	------	------------

No evidence from felt information of M greater than 4.

80.	1952	10	14	22	03	47.8	69.8	5.6 5.2
-----	------	----	----	----	----	------	------	------------

M_L from nearest station considered more representative.

81.	1953	01	24	09	58	49.1	66.0	5.3 4.6
-----	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 4 stations.

82.	1953	09	14	22	52	49.1	65.2	5.1 4.4
-----	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 3 stations.

83.	1954	01	10	21	04	49.2	68.2	3.9 3.1
-----	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 1 station

84.	1954	09	08	01	29	49.0	68.4	4.3 3.6
-----	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 1 station.

85.	1955	01	21	08	40	43.0	73.8	4.3 4.0
-----	------	----	----	----	----	------	------	------------

No evidence from felt information of M greater than 4.

86.	1955	02	03	02	30	44.5	73.2	4.3 4.0
-----	------	----	----	----	----	------	------	------------

No evidence from felt information of M greater than 4.

No.	Year	M	D	H	M	Lat.(°N)	Long.(°W)	Mag.
87.	1955	05	26	18	09	41.5	81.7	4.3 4.0
								Felt only locally.
88.	1955	08	16	07	35	42.9	78.3	4.3 4.0
								Felt only locally.
89.	1955	11	21	16	10	50.6	63.5	4.9 4.0
								$m_b(Lg)$ from 5 stations.
90.	1956	08	03	12	51	49.4	66.2	4.1 3.5
								$m_b(Lg)$ from 4 stations.
91.	1956	08	03	12	59	49.4	66.2	4.3 3.6
								$m_b(Lg)$ from 4 stations.
92.	1957	04	24	00	41	44.4	72.0	4.3 4.0
								Felt area consistent with M4 or smaller.
93.	1957	04	26	11	40	43.6	69.8	4.7 4.1
								$m_b(Lg)$ from 1 station.
94.	1957	10	16	19	13	50.5	64.9	4.8 4.1
								$m_b(Lg)$ from 6 stations.
95.	1958	05	14	17	41	47.0	76.6	5.4 5.0
								Felt area consistent with M5 or smaller.
96.	1958	08	08	22	15	47.9	70.4	4.0 3.6
								$m_b(Lg)$ from 1 station.

No.	Year	M	D	H	M	Lat.(°N)	Long.(°W)	Mag.
97.	1958	09	19	12	45	43.5	70.2	4.3 4.0

No evidence from felt information for M greater than 4.

98.	1961	07	05	22	43	50.3	66.7	5.0 4.3
-----	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 5 stations.

99.	1962	01	27	12	11	45.9	74.9	4.3 3.8
-----	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 1 station.

100.	1962	04	10	14	30	44.2	73.1	5.0 4.3
------	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 3 stations.

101.	1962	07	27	17	56	47.2	70.7	4.3 3.9
------	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 2 stations.

102.	1962	08	11	03	05	47.5	70.1	4.1 3.6
------	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 2 stations.

103.	1962	12	15	00	58	50.2	66.4	4.6 4.0
------	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 1 station.

104.	1963	10	15	12	29	46.2	77.6	4.4 4.1
------	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 2 stations.

105.	1963	10	15	13	59	46.2	77.6	4.5 4.2
------	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 2 stations.

106.	1965	10	05	14	36	49.8	67.7	4.6 3.9
------	------	----	----	----	----	------	------	------------

$m_b(Lg)$ from 5 stations.

